



Solution on Website:-

<https://physicsaholics.com/home/courseDetails/46>


Solution on YouTube:-

<https://youtu.be/veXEIzgr8t4>

- Q 1. Which of the following is true ?  
(a) Lyman series is a continuous spectrum  
(b) Paschen series is a line spectrum in the infrared  
(c) Balmer series is a line spectrum in the ultraviolet  
(d) The spectral series formula can be derived from the Rutherford model of the hydrogen atom
- Q 2. The wavelength of first line of Balmer series is  $6563 \text{ \AA}$ . The wavelength of first line of Lyman series will be  
(a)  $1215.4 \text{ \AA}$  (b)  $2500 \text{ \AA}$  (c)  $7500 \text{ \AA}$  (d)  $600 \text{ \AA}$
- Q 3. The wavelength of radiation required to excite an electron from first to third Bohr orbit in a doubly ionised lithium atom will be -  
(a)  $113.74 \text{ m}$  (b)  $113.74 \text{ cm}$   
(c)  $113.74 \text{ \AA}$  (d)  $113.74 \text{ mm}$
- Q 4. An excited hydrogen atom initially at rest in  $n = 3$  state, emits a photon by making a transition to ground to state. Then the momentum of the hydrogen atom will be (in N.s) -  
(a)  $6.45 \times 10^{-27}$  (b)  $6.63 \times 10^{-34}$   
(c)  $2.15 \times 10^{-27}$  (d) none of the above
- Q 5. When a hydrogen atom emits a photon of energy  $12.1 \text{ eV}$ , its orbital angular momentum changes by -  
(a)  $1.05 \times 10^{-34} \text{ J s}$   
(b)  $2.11 \times 10^{-34} \text{ J s}$   
(c)  $3.16 \times 10^{-34} \text{ J s}$   
(d)  $4.22 \times 10^{-34} \text{ J s}$
- Q.6 The ionization potential of H-atom is  $13.6 \text{ V}$ . The H-atoms in ground state are excited by mono chromatic radiations of photon energy  $12.09 \text{ eV}$ . Then the number of spectral lines emitted by the excited atoms, will be -  
(a) 1 (b) 2 (c) 3 (d) 4
- Q 7. Consider the spectral line resulting from the transition  $n = 2$  to  $n = 1$  in the atoms and ions given below, the shortest wavelength is produced by -  
(a) hydrogen atom  
(b) deuterium atom  
(c) singly ionized helium  
(d) doubly ionized lithium



- Q 8. Bohr's atom model assumes –
- (a) the nucleus is of infinite mass and is at rest
  - (b) electron in a quantized orbit will not radiate energy
  - (c) mass of the electron remains constant
  - (d) all of these
- Q 9. Figure represents in simplified form some of the energy levels of the hydrogen atom. The energy axis has a linear scale. If the transition of an electron from  $E_4$  to  $E_2$  were associated with the emission of blue light, which transition could be associated with the absorption of red light ?



- (a)  $E_4$  to  $E_1$
  - (b)  $E_3$  to  $E_2$
  - (c)  $E_2$  to  $E_3$
  - (d)  $E_1$  to  $E_4$
- Q 10. A mixture of ordinary hydrogen and tritium, is excited and its spectrum observed. Then, the ratio of the wavelengths of the  $H\alpha$  lines of the two kinds of hydrogen would be nearly -
- (a) 1 : 1
  - (b) 1 : 3
  - (c) 3 : 1
  - (d) nothing can be predicted
- Q 11. In hydrogen atom  $H\alpha$ -line arises due to transition  $n = 3 \rightarrow n = 2$ . In the spectrum of singly ionised helium there is a line having the same wavelength as the  $H\alpha$  line. This is due to the transition -
- (a)  $n = 3$  to  $n = 2$
  - (b)  $n = 2$  to  $n = 1$
  - (c)  $n = 5$  to  $n = 3$
  - (d)  $n = 6$  to  $n = 4$
- Q 12. Let  $n_1$  be the frequency of the series limit of the Lyman series,  $n_2$  be the frequency of the first line of the Lyman series, and  $n_3$  be the frequency of the series limit of the Balmer series -
- (a)  $n_1 - n_2 = n_3$
  - (b)  $n_2 - n_1 = n_3$
  - (c)  $n_3 = \frac{1}{2}(n_1 + n_2)$
  - (d)  $n_1 + n_2 = n_3$
- Q 13. Three photons coming from excited atomic-hydrogen sample are picked up. Their energies are 12.1 eV, 10.2 eV and 1.9 eV. These photons must come from -
- (a) a single atom
  - (b) two atoms
  - (c) three atoms
  - (d) either two atoms or three atoms



- Q 14. Radiations of wavelength  $\lambda$  are incident on hydrogen in the ground state. A fraction of these radiations absorbed by these atoms. There are ten different wavelength in the emission spectrum of excited atoms. The  $\lambda$  will be -
- (a)  $1211\text{\AA}$  (b)  $912\text{\AA}$   
(c)  $1211\text{\AA}$  (d)  $950.7\text{\AA}$
- Q 15. In which of the following transitions will the wavelength be minimum ?
- (a)  $n = 5$  to  $n = 4$  (b)  $n = 4$  to  $n = 3$   
(c)  $n = 3$  to  $n = 2$  (d)  $n = 2$  to  $n = 1$
- Q 16. If the wavelength of photon emitted due to transition of electron from third orbit to first orbit in a hydrogen atom is  $\lambda$ , then the wavelength of photon emitted due to of electron from fourth orbit to second orbit will be -
- (a)  $\frac{128}{27}\lambda$  (b)  $\frac{25}{9}\lambda$   
(c)  $\frac{36}{7}\lambda$  (d) None of these

PRATEEK JAIN  
PHYSICSAHOLICS

## Answer Key

Q.1 b	Q.2 a	Q.3 c	Q.4 a	Q.5 b
Q.6 c	Q.7 d	Q.8 d	Q.9 c	Q.10 a
Q.11 d	Q.12 a	Q.13 d	Q.14 d	Q.15 d
Q.16 a				