

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/88>

Video Solution on YouTube:-


[https://youtu.be/g69wcFX\\_VCE](https://youtu.be/g69wcFX_VCE)

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetalis/28>

- Q 1. We wish to observe a particle which is  $2.5 \text{ \AA}$  in size. The minimum energy photon that can be used (to observe a particle maximum required wavelength is equal to size of object) -
- (a) 5 KeV                      (b) 8 KeV                      (c) 10 KeV                      (d) 12 KeV

- Q 2. The radiation force experienced by body exposed to radiation of intensity  $I$  assuming surface of body to be perfectly absorbing is



- (a)  $\frac{\pi IR^2}{c}$   
 (b)  $\frac{2\pi IR^2}{c}$   
 (c)  $\frac{4\pi IR^2}{c}$   
 (d) None of these

PASSAGE (Q3. to Q5)

The light sensitive compound on most photographic films is silver bromide {AgBr}. A film is exposed when the light energy absorbed dissociates this molecule into its atoms, (the actual process is more complex, but the quantitative result does not differ with this assumption). The energy of dissociation of AgBr is  $10^5 \text{ J/mol}$ . Assume a photon that is just able to dissociate an AgBr molecule.

- Q 3. Find photon energy in e V.  
 (a) 1.04 eV                      (b) 1.14 eV                      (c) 1.72eV                      (d) 1.28 eV
- Q 4. Find the frequency of the photon  
 (a)  $2.71 \times 10^{14} \text{ Hz}$   
 (b)  $2.01 \times 10^{14} \text{ Hz}$   
 (c)  $2.5 \times 10^{14} \text{ Hz}$   
 (d)  $20.1 \times 10^{14} \text{ Hz}$
- Q 5. Will the radiation from a 50 kW, 100 MHz FM station expose the film?  
 (a) no                                      (b) yes  
 (c) cannot say                              (d) Incomplete data



- Q 6. A photon strikes a free electron at rest and is scattered straight backward. If the speed of electron after collision is  $\alpha c$ , where  $\alpha \ll 1$  then-
- (a) electron's kinetic energy is a fraction  $\alpha$  of photon's initial energy
  - (b) electron's kinetic energy is fraction  $\frac{1}{\alpha}$  of photon's initial energy
  - (c) electron's kinetic energy is a fraction  $\alpha^2$  of photon's initial energy
  - (d) electron's kinetic energy is a fraction  $\frac{1}{\alpha^2}$  of photon's initial energy
- Q 7. Two sources A and B have same power. The wavelength of radiation of A is  $\lambda_a$  and that of B is  $\lambda_b$ . The number of photons emitted per second by A and B are  $n_a$  and  $n_b$  respectively, then -
- (a)  $\lambda_a > \lambda_b$
  - (b) if  $\lambda_a > \lambda_b$ ,  $n_a < n_b$
  - (c) if  $\lambda_a < \lambda_b$ ,  $n_a < n_b$
  - (d) if  $\lambda_a > \lambda_b$ ,  $n_a = n_b$
- Q 8. A monochromatic beam of light ( $\lambda = 4900 \text{ \AA}$ ) incident normally upon a surface produces a pressure of  $5 \times 10^{-7} \text{ N/m}^2$  on it. Assuming that 25% of the light incident is reflected and the rest absorbed, find the number of photons falling per second on a unit area of this surface.
- (a)  $6 \times 10^{20} \text{ m}^{-2} \text{ s}^{-1}$
  - (b)  $2 \times 10^{20} \text{ m}^{-2} \text{ s}^{-1}$
  - (c)  $3 \times 10^{20} \text{ m}^{-2} \text{ s}^{-1}$
  - (d)  $9 \times 10^{20} \text{ m}^{-2} \text{ s}^{-1}$
- Q 9. If solar constant is S and average wavelength of sun radiation is  $\lambda$ , Total number of photons in a spherical volume of radius R near ground is
- (a)  $\frac{4\pi S \lambda R^3}{3hc^2}$
  - (b)  $\frac{2\pi S \lambda R^3}{3hc^2}$
  - (c)  $\frac{4\pi S \lambda R^3}{hc^2}$
  - (d) None of these
- Q 10. Two black objects of same diameter, a sphere and a disc, are placed in front of a uniform beam of light. The plane of disc is perpendicular to the light rays. The radiation force on them is
- (a) Zero
  - (b) Bigger on disc
  - (c) Bigger on sphere
  - (d) Same on both
- Q 11. A perfectly absorbing black solid sphere with constant density and radius R, hovers stationary above sun. This is because the gravitational attraction is balanced by radiation force due to sun light. Assume sun is far away that it is closely approximates a point source of light. The distance from the centre of sun at which sphere hovers is proportional to



- (a) R  
(c)  $\frac{1}{R^2}$

- (b)  $1/R$   
(d)  $R^0$

PRATEEK JAIN  
PHYSICSAHOLICS

## Answer Key

Q.1 a	Q.2 a	Q.3 a	Q.4 c	Q.5 a
Q.6 a	Q.7 c	Q.8 c	Q.9 a	Q.10 d
Q.11 d				