





- Q 7. A neutron moving with a speed  $v$  makes a head on collision with a hydrogen atom in ground state kept at rest. The minimum kinetic energy of neutron for which inelastic collision will take place is: (assume that mass of proton is nearly equal to the mass of neutron)
- (a) 10.2 eV                      (b) 20.4 eV                      (c) 12.1 eV                      (d) 16.8 eV
- Q 8. Hydrogen atoms absorb radiations of wavelength  $\lambda_0$  and consequently emit radiations of 6 different wavelengths of which three wavelengths are shorter than  $\lambda_0$ . Choose the correct alternative(s)
- (a) The final excited state of the atoms is  $n = 4$   
(b) The initial state of the atoms may be  $n = 2$   
(c) The initial state of the atoms may be  $n = 3$   
(d) There are three transitions belonging to Lyman series
- Q 9. In a hypothetical atom like that of hydrogen, the mass of the electrons is doubled. The energy  $E_0$  and radius  $r_0$  of the first Bohr orbit will be ( $a_0 =$  Bohr radius of hydrogen)
- (a)  $E_0 = -27.2 \text{ eV}$  ;  $r_0 = \frac{a_0}{2}$   
(b)  $E_0 = -27.2 \text{ eV}$  ;  $r_0 = a_0$   
(c)  $E_0 = -13.6 \text{ eV}$  ;  $r_0 = a_0/2$   
(d)  $E_0 = -13.6 \text{ eV}$  ;  $r_0 = \frac{a_0}{2}$
- Q 10. Suppose that the potential energy of an hypothetical atom consisting of a proton and an electron is given by  $U = -ke^2/3r^3$ . Then if Bohr's postulates are applied to this atom, then the radius of the  $n$ th orbit will be proportional to
- (a)  $n^2$                       (b)  $1/n^2$                       (c)  $n^3$                       (d)  $1/n^3$
- Q 11. Whenever a hydrogen atom emits a photon in the Balmer series,
- (a) it may emit another photon in the Balmer series  
(b) it must emit another photon in the Lyman series  
(c) the second photon will have a wavelength of about 122 nm  
(d) it may emit a second photon, but the wavelength of this photon cannot be predicted
- Q 12. A stationary  $\text{He}^+$  emitted a photon corresponding to the first line of Lyman series. This photon liberated a photoelectron from a stationary H-atom in the ground state. The velocity of the photoelectron will be –
- (a)  $3 \times 10^7 \text{ m/s}$                       (b)  $6 \times 10^6 \text{ m/s}$   
(c)  $8 \times 10^7 \text{ m/s}$                       (d)  $3.1 \times 10^6 \text{ m/s}$
- Q.13 An electron of energy 10.8 eV undergoes an inelastic collision with a hydrogen atom in its ground state. Then (assuming  $m_H \gg m_e$ , neglecting recoil of atom) -
- (a) the outgoing electron has energy 10.8 eV  
(b) 10.2 eV of the incident electrons energy is absorbed by H-atom and the electron would come out with 0.6 eV energy  
(c) the entire energy is absorbed by H-atom and the electron stops  
(d) none of the above



- Q.14 If we take into account the reality that both the nucleus and electron revolve around their common centre of mass. During electron transition from a higher state  $n_2$ , to a lower state,  $n_1$ , we find that the wavelength of the photon emitted is not given by the formula  $\frac{1}{\lambda} = R \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$  where R is the Rydberg constant. The correct wavelength, in that case depends on mass of electron (m) and mass of the nucleus (M) and is given by
- (a)  $\frac{1}{\lambda} = R \frac{m}{M} \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$   
(b)  $\frac{1}{\lambda} = R \left( 1 + \frac{m}{M} \right) \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$   
(c)  $\frac{1}{\lambda} = R \left( \frac{m}{n_1^2} - \frac{M}{n_2^2} \right)$   
(d)  $\frac{1}{\lambda} = R \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \left( \frac{M}{M+m} \right)$
- Q.15 An H atom in ground state with kinetic energy 22eV hits another stationary H-atom in ground state. The collision:
- (a) must be elastic  
(b) may be elastic  
(c) may be perfectly inelastic  
(d) may be inelastic
- Q.16 Wavelength of photon emitted by H atom in  $n = 4$  to  $n = 2$  transition is equal to wavelength of photon produced by He atom in
- (a)  $n = 4$  to  $n = 2$  transition  
(b)  $n = 2$  to  $n = 1$  transition  
(c)  $n = 6$  to  $n = 3$  transition  
(d)  $n = 8$  to  $n = 4$  transition

## Answer Key

<b>Q.1</b>	<b>a</b>	<b>Q.2</b>	<b>a</b>	<b>Q.3</b>	<b>c</b>	<b>Q.4</b>	<b>a</b>	<b>Q.5</b>	<b>a</b>
<b>Q.6</b>	<b>b</b>	<b>Q.7</b>	<b>b</b>	<b>Q.8</b>	<b>a,b,d</b>	<b>Q.9</b>	<b>a</b>	<b>Q.10</b>	<b>b</b>
<b>Q.11</b>	<b>b,c</b>	<b>Q.12</b>	<b>d</b>	<b>Q.13</b>	<b>b</b>	<b>Q.14</b>	<b>d</b>	<b>Q.15</b>	<b>b,d</b>
<b>Q.16</b>	<b>d</b>								