

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/42>


Video Solution on YouTube:-

<https://youtu.be/rTg2MIwPV6g>

Written Solution on Website:-


<https://physicsaholics.com/note/notesDetailis/36>

- Q 1. If the velocity of block B in the given arrangement is 300 mm/sec towards right. Find the velocity of A:




- (a) 100 mm/sec (b) 200 mm/sec
(c) 300 mm/sec (d) 400 mm/sec

- Q 2. Find the velocity of block B when ring A is moving downward with velocity v :




- (a) $v \sin \theta$ (b) $\frac{v}{2} \sin \theta$
(c) $v \cos \theta$ (d) $\frac{v}{2} \cos \theta$

- Q 3. If block A is moving horizontally with velocity V_A , then find the velocity of block B at the instant as shown in fig:.




- (a) $\frac{hV_A}{2\sqrt{x^2+h^2}}$ (b) $\frac{xV_A}{\sqrt{x^2+h^2}}$
 (c) $\frac{xV_A}{2\sqrt{x^2+h^2}}$ (d) $\frac{hV_A}{\sqrt{x^2+h^2}}$

- Q 4. A cart is being pulled up the incline, using a motor M and an ideal pulley and ideal rope arrangement as shown in figure. Then the speed of point ' P ' of the string with which it moves so that the car moves up the inclined plane with a constant speed of $V_{cart} = 2 \text{ m/s}$ is (Incline is at rest):




- (a) 12 m/s (b) 3 m/s
 (c) 5 m/s (d) 6 m/s

- Q 5. In Fig. a ball of mass m_1 and a block of mass m_2 are joined together with an inextensible string. The ball can slide on a smooth horizontal surface. If V_1 and V_2 are the respective speeds of the ball and the block, then determine the constraint relation between velocities of the two.




- (a) $V_2 = V_1 \cos \theta$ (b) $V_1 = V_2 \cos \theta$
 (c) $V_1 = V_2$ (d) $V_2 = V_1 \sin \theta$

Q 6. Find $V_B = ?$




- (a) 10 m/s (b) 8 m/s
(c) 14 m/s (d) 6 m/s

Q 7. Determine the speed with which block B rises in Fig. if the end of the cord at A is pulled down with a speed of 2 m/s.




- (a) 4 m/s (b) 3 m/s
(c) $\frac{3}{2}$ m/s (d) $\frac{1}{2}$ m/s

Q 8. Two rings each of mass $M = 100 \text{ gm}$ are constrained to move along a fixed horizontal rod. An ideal string is connected with rings and block of mass $M_0 = 200 \text{ gm}$ is connected to the mid point of string. At a certain moment the mass m is moving downward with velocity $\sqrt{3} \text{ m/s}$. Find the speed of ring of M at the moment:




- (a) 4 m/s (b) 3 m/s
(c) 2 m/s (d) 1 m/s

Q 9. In the given figure, find the speed of pulley P –




- (a) $\frac{V}{2}$ (b) $2V \cos \theta$
 (c) $\frac{2V}{\cos \theta}$ (d) $\frac{V}{2 \sin \theta}$

Q 10. Figure shows a rod of length l resting on a wall and the floor. Its lower end A is pulled towards left with a constant velocity u . As a result of this, end A starts moving down along the wall. Find the velocity of the other end B downward when the rod makes an angle θ with the horizontal:



- (a) $u \tan \theta$ (b) $u \cot \theta$
 (c) $u \sin \theta$ (d) $u \cos \theta$

Q 11. The velocities of A and B are marked in the figure. Find the velocity of block C (assume that the pulleys are ideal and string inextensible)



- (a) 2 m/s (b) 4 m/s
 (c) 5 m/s (d) $\sqrt{10}$ m/s



Answer Key

Q.1 b	Q.2 c	Q.3 c	Q.4 d	Q.5 a
Q.6 a	Q.7 d	Q.8 d	Q.9 c	Q.10 b
Q.11 c				

PRATEEK JAIN
PHYSICSAHOLICS