



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/46>

Video Solution on YouTube:-

https://youtu.be/81zl6Y_7sTc

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/19>

- Q 1. One electron & one proton is accelerated by equal potential. Ratio in their de-broglie wavelength is-
- (a) $\sqrt{\frac{m_p}{m_e}}$ (b) $\frac{m_e}{m_p}$
(c) $\frac{m_p}{m_e}$ (d) 1
- Q 2. The de-Broglie wavelength associated with an electron having a kinetic energy of 10 eV is
(a) 10 Å (b) 12.27 Å (c) 3.9 Å (d) 0.10 Å
- Q 3. A double slit interference experiment is performed by a beam of electrons of energy 100 eV and the fringe spacing is observed to be β . Now if the electrons energy is increased to 10 keV, then the fringe spacing -
(a) remains the same (b) becomes 10β
(c) becomes 100β (d) becomes $\beta/10$
- Q 4. If E_1 , E_2 and E_3 are the respective kinetic energies of an electron, an alpha particle and a proton, each having the same de Broglie wavelength, then
(a) $E_1 > E_3 > E_2$ (b) $E_2 > E_3 > E_1$
(c) $E_1 > E_2 > E_3$ (d) $E_1 = E_2 = E_3$
- Q 5. If the momentum of electron is changed by P_m then the De Broglie wavelength associated with it changes by 0.50 %. The initial momentum of electron will be -
(a) $\frac{P_m}{200}$ (b) $\frac{P_m}{100}$ (c) 200 Pm (d) 400 Pm
- Q.6 The thermal energy of a particle at temperature T°K is kT, then the associated de-Broglie wavelength will be -
(a) h/mkT (b) $\frac{h}{\sqrt{2mkT}}$
(b) $\frac{h}{2mkT}$ (d) $\frac{2h}{mkT}$
- Q 7. Wrong statement in connection with Davisson-Germer experiment is -
(a) The inter-atomic distance in nickel crystal is of the order of the de-Broglie wavelength
(b) Electrons of constant energy are obtained by the electron gun
(c) Nickel crystal acts as a three-dimensional diffracting grating
(d) Davission-Germer experiment is a photoelectric experiment



- Q 8. In Davisson-Germer experiment the relation between scattering angle θ and glancing angle ϕ is -
- (a) $\theta = 90^\circ - \phi$ (b) $\theta = \frac{90^\circ - \phi}{2}$
(c) $\theta = 180^\circ - \phi$ (d) $\phi = \left(\frac{180^\circ - \theta}{2}\right)$
- Q 9. The de-Broglie wavelength of a vehicle moving with velocity v is λ . Its load is changed so that the velocity as well as the kinetic energy are doubled. Then the new de-Broglie wavelength of the vehicle will be -
- (a) λ (b) 2λ (c) $\lambda/2$ (d) $\lambda/4$
- Q 10. An electron is confined to a tube of length L . The electron's potential energy in one half of the tube is zero, while the potential energy in the other half is 10 eV. If the electron has a total energy $E = 15$ eV, then the ratio of the de-Broglie wavelength of the electron in the 10 eV region of the tube to that in the other half is -
- (a) $1/\sqrt{3}$ (b) $\sqrt{3}$ (c) 3 (d) $\frac{1}{3}$
- Q 11. In majority of crystals the value of lattice constant is of the order of 3\AA . The proper X-rays with which the crystal structure can be studied are -
- (a) 50\AA to 100\AA (b) 10\AA to 50\AA
(c) 5\AA to 10\AA (d) 0.1\AA to 2.7\AA

Answer Key

Q.1 a	Q.2 c	Q.3 d	Q.4 a	Q.5 c
Q.6 b	Q.7 d	Q.8 d	Q.9 a	Q.10 b
Q.11 d				