



DPP – 1 (Sound Wave)

Video Solution on Website:-

https://physicsaholics.com/home/courseDetails/94

Video Solution on YouTube:-

https://youtu.be/rhMRmNgKw 8

Written Solution on Website:-

https://physicsaholics.com/note/notesDetalis/45


Q 1. **State True or false**

 S_1 : If pressure of a gas is increased keeping the temperature constant the speed of sound wave in the gas increases.

 S_2 : In a travelling sinusoidal sound wave the phase difference between displacement wave and pressure wave is $\frac{\pi}{2}$.

 S_3 : At same temperature sound propagates with same speed in CO gas and N_2 gas (a) T F T

Figure shown is a graph, at a certain time t, of the displacement function S(x,t) of three sound Q 2. waves 1,2 and 3 as marked on the curves that travel along x-axis through air, If P_1 , P_2 and P_3 represent their pressure amplitudes respectively, then correct relation between them is:




(b) $P_3 > P_2$

(c) $P_1 = P_2 = P_3$

03. The maximum pressure variation that the human ear can tolerate in loud sound is about 30 N/m². The corresponding maximum displacement amplitude for a sound wave in air having a frequency of 10^4 Hz is:(take velocity of sound in air as 300 m/s and density of air 1.5 kg/m³) (a) $\frac{2\pi}{3} \times 10^{-2}$ (b) $\frac{2\times10^{-4}}{\pi}$ (c) $\frac{\pi}{3} \times 10^{-2}$ (d) $\frac{10^{-4}}{30\pi}$

The graph below shows displacement(y) versus distance (x) along the line of a transmission Q 4. of a longitudinal wave at a given instant. The positive y-axis is displacement to the right and the negative y-axis is displacement to the left. The pressure maxima positions happen at the positions marked -



(a) P

(b) Q

(c) R

(d) S



P hysicsaholics



Q 5.	minimum pressures at a		$^{-3}$ N/ m^2 , find the amp ir 1.4×10^5 N/ m^2 .	etween the maximum and blitude of vibration of the $ (d) \ 2.2 \times 10^{-10} \ m. $		
Q 6.		in human mind remains for 0.1 second. A man is clapping near a wall. e of wall from man to listen echo of clap? Velocity of sound in air is				
	(a) 17m	(b) 16m	(c) 16.5m	(d) 17.5m		
Q 7.	A to B. Absolute tempe	wo points A and B are separated by distance 960 m. Air temperature is varying linearly from to B. Absolute temperature of B is 4 times that of A. Velocity of sound at A is 320m/sec. A bund is produced at A, Time taken by it to reach B is 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1				
Q 8.		waves in an ideal gas at the ty of gas molecules at the (b) $\frac{v_1}{v_2} = \sqrt{\gamma} \frac{\omega_1}{\omega_2}$ (c) $\frac{v_1}{v_2}$	ese two temperatures			
Q 9.	Of the following properties of a sound wave, the one that is independent of the other is its - (a) Amplitude (b) Velocity (c) Wavelength (d) Frequency					
Q 10.	 Mark the correct statement (a) According to Newtons formula velocity of sound in hydrogen is 4 times that in oxygen at same temperature. (b) According to Laplace formula velocity of sound in hydrogen is 4 times that in oxygen at same temperature. (c) According to Newtons formula velocity of sound in helium is 2√2 times that in oxygen at same temperature. (d) According to Laplace formula velocity of sound in helium is more than 2√2 times that in oxygen at same temperature. 					
Q 11.	Velocity of sound in air is (a) Smaller at night due to low temperature at night. (b) Greater at night due to low temperature at night (c) Smaller at night due to low density of air at night. (d) Greater at night due to low density of air at night.					
Q 12.	Two sound waves are moving in just opposite directions, first wave due east and second wave due west. At a given time instant a particle of first wave is at its one extreme position which is east of its mean position and at same time instant a particle of second wave is at its one extreme position which is west of its mean position. Phase difference between pressure variation at these two particles is (a) 0 (b) π (c) $\frac{\pi}{2}$ (d) None of these					
Q 13.	Colum	and in gas match the follown-I rature of gas is	Col	lumn-II ed becomes		



Physicsaholics



made 4 times and pressure 2 times

(B) Only pressure is made 4 time without change in temperature

(C) Only temperature is changed to 4 times

(D) Molecular mass of the gas is made 4 times

 $2\sqrt{2}$ times

(Q) speed becomes 2 times

(R) speed remains unchanged

(S) speed remains half



Answer Key

Q.1 b	Q.2 b	Q.3 d	Q.4 c	Q.5 d
Q.6 c	Q.7 d	Q.8 a	Q.9 b	Q.10 a, b, c, d
Q.11 d	Q.12 a			

Q.13 A \rightarrow Q; B \rightarrow R; C \rightarrow Q; D \rightarrow S