

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/38>


Video Solution on YouTube:-

<https://youtu.be/pqSwR5H6gMY>

Written Solution on Website:-


<https://physicsaholics.com/note/notesDetails/43>

- Q 1. A body slides down a frictionless track which ends in a circular loop of diameter  $D$ , then the minimum height  $h$  of the body in term of  $D$  so that it may just complete the loop, is
- (a)  $h = \frac{5D}{2}$  (b)  $h = \frac{5D}{4}$   
 (c)  $h = \frac{3D}{4}$  (d)  $h = \frac{D}{4}$
- Q 2. A car moving with speed  $30 \text{ m/s}$  on a circular path of radius  $500\text{m}$ . Its speed is increasing at the rate of  $2 \text{ m/s}^2$ . The acceleration of the car is
- (a)  $9.8 \text{ m/s}^2$  (b)  $1.8 \text{ m/s}^2$   
 (c)  $2 \text{ m/s}^2$  (d)  $2.7 \text{ m/s}^2$
- Q 3. A stone tied to the end of a string which is  $80\text{cm}$  long is whirled in a horizontal circle with a constant speed. If the stone makes  $14$  revolutions in  $25\text{s}$ , Find work done by tension on stone
- (a)  $9.91 \text{ J}$  (b)  $14 \text{ J}$   
 (c)  $12.69 \text{ J}$  (d) zero
- Q 4. The string of pendulum of length  $l$  is displaced through  $90^\circ$  from the vertical and released. Then the minimum strength of the string in order to withstand the tension, as the pendulum passes through the mean position is
- (a)  $mg$  (b)  $3mg$   
 (c)  $5mg$  (d)  $6mg$
- Q 5. A frictionless track ABCDE ends in a circular loop of radius  $R$ . A body slides down the track from point A which is at height  $h = 5\text{cm}$ . Maximum value of  $R$  for a body to complete the loop successfully is




- (a) 2 cm                      (b)  $\frac{10}{3}$  cm  
 (c)  $\frac{15}{4}$  cm                  (d)  $\frac{18}{3}$  cm

Q 6. A cube of mass  $M$  starts from rest from point 1 at a height  $4R$ , where  $R$  is the radius of the circular track. The cube slides down on the frictionless track and around the loop. The force which the track exerts on the cube at point 2 is:



- (a)  $mg$                                       (b)  $2mg$   
 (c)  $3mg$                                       (d)  $4mg$

Q 7. A block shown in figure slides on a semicircular frictionless track. If it starts from rest at position A, what is its speed at the point marked B? Take  $g=10\text{m/s}^2$




- (a) 2.21 m/s                                      (b) 1.32 m/s  
 (c) 7.54 m/s                                      (d) 3.76 m/s

Q 8. A ball is released from point A as shown in figure. The ball leaves the track at B. All surfaces are smooth. If track makes an angle  $30^\circ$  with horizontal at B, then maximum height attained by ball will be




- (a) 4m    (b) 2m  
 (c) 3m    (d) 1m

Q 9. A particle originally at rest at the highest point of a smooth vertical circle is slightly displaced. It will leave the circle at a vertical distance  $h$  below the highest point such that  $h = ?$




- (a)  $R$                       (b)  $\frac{R}{3}$   
 (c)  $\frac{2R}{3}$                     (d)  $\frac{R}{2}$

Q 10. A small particle of mass  $m$  attached with a light inextensible thread of length  $L$  is moving in a vertical circle. In the given case particle is moving in complete vertical circle and ratio of its maximum to minimum velocity is 2:1. Velocity of the particle when it is moving vertically downward is



- (a)  $\sqrt{\frac{10gL}{3}}$                       (b)  $2\sqrt{\frac{gL}{3}}$   
 (c)  $\sqrt{\frac{8gL}{3}}$                         (d)  $\sqrt{\frac{13gL}{3}}$

Q 11. A particle is given an initial speed  $u$  inside a smooth spherical shell of radius  $R=1$  m such that it is just able to complete the circle. Acceleration of the particle when its velocity is vertical is




- (a)  $g\sqrt{10}$                       (b)  $g$   
 (c)  $g\sqrt{2}$                         (d)  $3g$

Q 12. A pendulum bob on a 2 m string is displaced  $60^\circ$  from the vertical and then released. What is the speed of the bob as it passes through the lowest point in its path



- (a)  $\sqrt{2}$  m/s                      (b)  $\sqrt{9.8}$  m/s  
 (c) 4.43 m/s                        (d)  $\frac{1}{\sqrt{2}}$  m/s

Q 13. A particle of mass  $m$  is attached to one end of a light inextensible string and the other end of the string is fixed in vertical plane as shown. Particle is given a horizontal velocity  $u = \sqrt{\frac{5}{2}gl}$ . The maximum angle made by string with downward vertical is



- (a)  $\cos^{-1}\left(\frac{1}{4}\right)$                       (b)  $\sin^{-1}\left(\frac{1}{4}\right)$   
 (c)  $\frac{\pi}{2} + \sin^{-1}\left(\frac{1}{4}\right)$                       (d)  $\frac{\pi}{2} - \sin^{-1}\left(\frac{1}{4}\right)$

PRATEEK JAIN  
PHYSICSAHOLICS

## Answer Key

Q.1 b	Q.2 d	Q.3 d	Q.4 b	Q.5 a
Q.6 c	Q.7 d	Q.8 c	Q.9 b	Q.10 a
Q.11 a	Q.12 c	Q.13 c		