



## **DPP – 4 (Wave Optics)**

Video Solution on Website:-

https://physicsaholics.com/home/courseDetails/33

Video Solution on YouTube:-

https://youtu.be/uzJxr7xhyGY

Written Solution on Website:-

https://physicsaholics.com/note/notesDetalis/46

- Q 1. Unpolarized light is incident on a plane glass surface. What should be the angle of incidence so that the reflected and refracted rays are perpendicular to each other? (Given refractive index of glass, n=1.5)
  - (a)  $\sin^{-1} 1.5$
- (b)  $tan^{-1} 1.5$
- (c)  $\cos^{-1} 1.5$
- (d)  $\sin^{-1}\frac{2}{3}$
- Q 2. Light waves can be polarized as they are
  - (a) Transverse
- (b) Of high frequency
- (c) Longitudinal
- (d) Reflected
- In the case of linearly polarized light, the magnitude of the electric field vector O 3.
  - (a) Is parallel to the direction of propagation
  - (b) Does not change with time
  - (c) Increases linearly with time
  - (d) Varies periodically with time
- The angle of polarization for any medium is 60°, what will be critical angle for this Q 4.

(a) 
$$\sin^{-1} \sqrt{3}$$

(b) 
$$\tan^{-1}\sqrt{3}$$

(c) 
$$\cos^{-1}\sqrt{3}$$

(d) 
$$\sin^{-1} \frac{1}{\sqrt{3}}$$

- Q 5. A polaroid is placed at  $45^{\circ}$  to an incoming light of intensity  $I_0$ . Now the intensity of light passing through polaroid after polarization would be
- (b)  $\frac{I_0}{2}$
- $(c)\frac{I_0}{4}$
- (d) zero
- Unpolarized light falls on two polarizing sheets placed one on top of the other. What O 6. must be the angle between the characteristic directions of the sheets if the intensity of the transmitted light is one third of intensity of the incident beam?
  - (a)  $\tan^{-1}\left(\sqrt{\frac{3}{2}}\right)$

(b)  $\sin^{-1}\left(\sqrt{\frac{3}{2}}\right)$ (d)  $\sin^{-1}\left(\sqrt{\frac{1}{3}}\right)$ 

(c)  $\cos^{-1} \left( \frac{1}{2} \right)^{\frac{2}{3}}$ 

- Q 7. Unpolarized light of intensity I<sub>0</sub> is incident on a polarizer and the emerging light strikes a second polarizing filter with its axis at 45° to that of the first. The intensity of the emerging beam



## hysicsaholics



(a)  $\frac{I_0}{2}$ 

(b)  $\frac{I_0}{4}$ 

(c)  $I_0$ 

(d)  $\frac{I_0}{3}$ 

Q 8. A beam of the plane polarized light having flux  $10^{-3}$  watt falls normally on a polarizer of a cross sectional area  $3 \times 10^{-4} \ m^2$ . The polarizer rotates with an angular frequency of 31.4 rad/s. The energy of the light passes through the polarizer per revolution will be

(a)  $10^{-4}$  J

(b)  $10^{-3}$  J

 $(c) 2 \times 10^{-4} J$ 

(d)  $3 \times 10^{-4} \text{ J}$ 

Q 9. The angle between pass axis of polarizer and analyzer is 45°. The percentage of polarized light passing through analyzer is

(a) 100 %

(b) 50 %

(c) 25 %

(d) 75 %

Q 10. A ray of light is incident on the surface of a glass plate at an angle of incidence equal to Brewster's angle  $\theta$ . If  $\mu$  represents the refractive index of glass with respect to air, then the angle between reflected and refracted rays is

(a)  $90^{\circ} + \theta$ 

(b)  $\sin^{-1}(\mu\cos\theta)$ 

(c) 90°

(d)  $90^{\circ} - \sin^{-1} \left( \sin \frac{\theta}{u} \right)$ 

- Q 11. Polarized glass is used in sun glasses because
  - (a) It reduces the light intensity to half an account of polarization
  - (b) It is fashionable
  - (c) It has good colour
  - (d) It is cheaper
- Q 12. A light has amplitude A (after polarizer) and angle between analyzer and polarizer is 60°. Light is reflected by analyzer has amplitude

(a)  $A\sqrt{2}$ 

(b)  $\frac{A}{\sqrt{2}}$ 

(c)  $\frac{\sqrt{3}A}{2}$ 

 $(d)\frac{A}{2}$ 

- Q 13. When a plane polarized light is passed through an analyzer and analyzer is rotated through 90°, the intensity of the emerging light
  - (a) Varies between maximum and zero
  - (b) Becomes zero
  - (c) Does not vary
  - (d) cant say anything
- Q 14. When the angle of incidence on a material is 60°, the reflected light is completely polarized. The velocity of the refracted ray inside the material is (in m/s)

(a)  $3 \times 10^8$ 

(b)  $\frac{3}{\sqrt{2}} \times 10^8$ 

(c)  $\sqrt{3} \times 10^{8}$ 


(d)  $0.5 \times 10^8$ 




## Physicsaholics



Q 15. Unpolarized light beam of intensity  $I_0$  is incident on polaroid  $P_1$ . The three polaroids are arranged in such a way that transmission axis of  $P_1$  and  $P_3$  are perpendicular to each other. Angle between the transmission axis of  $P_2$  and  $P_3$  is  $60^\circ$ . The intensity of the beam coming out from  $P_3$  will be



- (a)  $\frac{I_0}{2}$
- (b)  $\frac{3I_0}{8}$
- $\frac{3I_0}{32}$  (d)  $\frac{3}{6}$



Q.1 b	Q.2 a	Q.3 d	Q.4 d	Q.5 b
Q.6 c	Q.7 c	Q.8 a	Q.9 b	Q.10 c
Q.11 a	Q.12 d	Q.13 a	Q.14 c	Q.15 c