

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/37>


Video Solution on YouTube:-

<https://youtu.be/VkqGHJ8gU3E>


Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/49>

- Q 1. A smooth sphere is moving on a horizontal surface with velocity vector $2\hat{i} + 2\hat{j}$ immediately before it hits a vertical wall. The wall is parallel to \hat{j} vector and the coefficient of restitution between the sphere and the wall is $e = \frac{1}{2}$. The velocity vector of the sphere after it hits the wall is:
- (a) $\hat{i} - \hat{j}$ (b) $-\hat{i} + 2\hat{j}$
 (c) $-\hat{i} - \hat{j}$ (d) $2\hat{i} - \hat{j}$
- Q 2. A sphere has a elastic oblique collision with another identical sphere which is initially at rest. The angle between their velocities after the collision is
- (a) 30° (b) 45°
 (c) 60° (d) 90°
- Q 3. A ball of mass m moving horizontally with velocity u hits a wedge of mass M . The wedge is situated on a smooth horizontal surface. If after striking with wedge the ball starts moving in vertical direction and the wedge starts moving in horizontal plane. Calculate the velocity of wedge V




- (a) $\frac{mu}{M}$ (b) $\frac{mu(\sin \theta - 1)}{M \sin \theta}$
 (c) $\frac{mu(\cos \theta - 1)}{M \sin \theta}$ (d) $\frac{mu(\sin 2\theta - 1)}{M \cos \theta}$
- Q 4. A ball of mass m moving horizontally with velocity u hits a wedge of mass M . The wedge is situated on a smooth horizontal surface. If after striking with wedge the ball starts moving in vertical direction and the wedge starts moving in horizontal plane. The impulse imparted by the ball on the wedge




- (a) $\frac{mu}{\sin \theta}$ (b) $\frac{mu}{\cos \theta}$
 (c) $mu \sin \theta$ (d) $\frac{mu}{\cot \theta}$

Q 5. A ball of mass m moving horizontally with velocity u hits a wedge of mass M . The wedge is situated on a smooth horizontal surface. If after striking with wedge the ball starts moving in vertical direction and the wedge starts moving in horizontal plane. The coefficient of restitution $e = ?$



- (a) $\frac{m}{M} \cot^2 \theta$ (b) $\frac{m}{M} \tan^2 \theta$
 (c) $\frac{m}{M} + \cot^2 \theta$ (d) $\frac{m}{M} - \cot^2 \theta$

Q 6. A ball of mass m collides with a stationary wedge of mass M , perpendicular to its inclined face, inclined at an angle as shown in the figure. If the coefficient of restitution between the wedge and ball is e , calculate the velocity of wedge just after collision.




- (a) $\frac{mu \sin \alpha (e+2)}{M+m \sin^2 \alpha}$ (b) $\frac{mu \sin \alpha (e+1)}{M+m \sin^2 \alpha}$
 (c) $\frac{mu \sin \alpha (e+1)}{2M+m \sin^2 \alpha}$ (d) $\frac{mu \sin \alpha (e+2)}{M+2m \sin^2 \alpha}$

Q 7. In a collision between two solid spheres, velocity of separation along the line of impact (assume no external forces act on the system of two spheres during impact)

- (a) cannot be greater than velocity of approach
 (b) cannot be less than velocity of approach


- (c) cannot be equal to velocity of approach
 (d) none of these

Q 8. A ball of mass m moving vertically down, collides with inclined surface of the wedge. After the collision, wedge starts moving in horizontal direction with velocity v_0 . If all the surfaces are smooth then impulse applied by wedge on the ball during collision is given by



- (a) $mv_0 \sin \theta$ (b) $mv_0 \cos \theta$
 (c) $\frac{mv_0}{\sin \theta}$ (d) $\frac{mv_0}{\cos \theta}$

Q 9. A ball of mass $m = 1\text{kg}$ falling vertically with a velocity $v_0 = 2\text{m/s}$ strikes a wedge of mass $M = 2\text{kg}$ kept on a smooth, horizontal surface as shown in figure. If impulse between ball and wedge during collision is J . Find impulse on wedges from ground during impact.



- (a) $\frac{1}{2} J$ (b) $\frac{\sqrt{3}}{2} J$
 (c) $\frac{1}{\sqrt{3}} J$ (d) $\frac{2}{\sqrt{3}} J$

Q 10. A sphere of mass $m_1 = 2\text{kg}$ collides with a sphere of mass $m_2 = 3\text{kg}$ which is at rest. Mass m_1 will move at right angles to the line, joining centers at the time of collision, if the coefficient of restitution is

- (a) $\frac{4}{9}$ (b) $\frac{1}{2}$
 (c) $\frac{2}{3}$ (d) $\sqrt{2/3}$

Q 11. Sand is being dropped on a conveyer belt at the rate of $M \text{ kg/s}$. The force (in newton) necessary to keep the belt moving with a constant velocity of $v \text{ m/s}$ will be

- (a) $\frac{Mv}{2}$ (b) zero
 (c) Mv (d) $2Mv$



Answer Key

Q.1 b	Q.2 d	Q.3 a	Q.4 a	Q.5 c
Q.6 b	Q.7 a	Q.8 c	Q.9 b	Q.10 c
Q.11 c	Q.12 c	Q.13 a	Q.14 d	Q.15 a

PRATEEK JAIN
PHYSICSAHOLICS