

DPP – 2 (Wave Optics)

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/96>


Video Solution on YouTube:-

<https://youtu.be/7AM7-YXYfYE>


Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/47>

- Q 1. Two identical narrow slits S_1 and S_2 are illuminated by light of wavelength λ from a point source P. If, as shown in the diagram above the light is then allowed to fall on a screen, and if n is a positive integer the condition for destructive interference at Q is that



- (a) $(l_1 - l_2) = (2n + 1)\lambda/2$
 (b) $(l_3 - l_4) = (2n + 1)\lambda/2$
 (c) $(l_1 + l_2) - (l_2 + l_4) = n\lambda$
 (d) $(l_1 + l_3) - (l_2 + l_4) = (2n + 1)\lambda/2$
- Q 2. For maxima (bright fringe) at point P, relation between given quantities is (angles shown in figure are not small)




- (a) $|d \sin \phi - d \sin \theta| = (2n-1)\lambda/2$
 (b) $|d \sin \phi - d \sin \theta| = n\lambda$
 (c) $|d \sin \phi - d \sin \theta| = (2n-1)\lambda/4$
 (d) None of these
- Q 3. Two coherent point sources s_1 and s_2 vibrating in phase emit light of wavelength λ . The separation between the sources is 2λ . The smallest distance from s_2 on a line passing through s_2 and perpendicular to s_1s_2 where a minimum of intensity occurs is:
- (a) $\frac{7\lambda}{12}$ (b) $\frac{15\lambda}{4}$ (c) $\frac{\lambda}{2}$ (d) $\frac{3\lambda}{4}$
- Q 4. White light is used to illuminate the two slits in Young's double slit experiment. The separation between the slits is b and the screen is at a distance d ($\gg b$) from the slits. At a point on the screen directly in front of one of the slits, certain wavelengths are missing. Some of these missing wavelengths are:
- (a) $X = b^2/d$ (b) $\lambda = 2b^2/d$ (c) $\lambda = b^2/3d$ (d) $\lambda = 2b^2/3d$




- Q 5. In a Biprism experiment, if the wavelength of red light used is 6.5×10^{-7} m and that of green is 5.2×10^{-7} m, the value of n for which $(n + 1)$ th green bright band coincides with the n th red bright band for the same setting is given by -
 (a) 2 (b) 3 (c) 4 (d) 1


- Q 6. In a YDSE experiment if a slab whose refractive index can be varied is placed in front of one of the slits then the variation of resultant intensity at mid-point of screen with ' μ ' will be best represented by ($\mu \geq 1$). [Assume slits of equal width and there is no absorption by slab]




(a)



(b)




(c)



(d)

- Q 7. If white light is used in a Young's double-slit experiment -
 (a) bright white fringe is formed at the centre of the screen
 (b) fringes of different colours are observed clearly only in the first order
 (c) the first-order violet fringes are closer to the centre of the screen than the first order red fringes
 (d) the first-order red fringes are closer to the centre of the screen than the first order violet fringes

- Q 8. A parallel beam of light ($\lambda = 5000 \text{ \AA}$) is incident at an angle $\alpha = 30^\circ$ with the normal to the slit plane in a young's double slit experiment. Assume that the intensity due to each slit at any point on the screen is I_0 . Point O is equidistant from S_1 & S_2 . The distance between slits is 1mm.



- (a) the intensity at O is $4I_0$
 (b) the intensity at O is zero
 (c) the intensity at a point on the screen 4m from O is $4I_0$
 (d) the intensity at a point on the screen 4m from O is zero
- Q 9. Two coherent narrow slits S_1 and S_2 emitting light of wavelength λ in the same phase are placed parallel to each other at a small separation of 3λ . The light is collected on a screen S which is placed at a distance D ($\gg \lambda$) from the slit S_1 and shown in figure. Find the distance x such that the intensity at point P is equal to the intensity at O.

