



## DPP – 3 (Wave Optics)

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/96>

Video Solution on YouTube:-


<https://youtu.be/6hsZg25muOo>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/47>

- Q 1. The minimum thickness of film which will strongly reflect the light of  $\lambda = 300$  nm, the R.I of material of film is 1.25  
(a) 120 nm                      (b) 100 nm                      (c) 110 nm                      (d) 60 nm

- Q 2. A thin slice is cut out of a glass cylinder along a plane parallel to its axis. The slice is placed on a flat plate as shown. The observed interference fringes from this combination shall be:




- (a) straight  
(b) circular  
(c) equally spaced  
(d) having fringe spacing which increases as we go outwards
- Q 3. A thin film of index 1.6 is placed in air and white light reflected from it is viewed. The wavelengths 432 nm, 540 nm and 120 nm are missing. Then -  
(a) Minimum possible thickness of film is 675 nm  
(b) if the order of 432 nm is 5, the order of 540 nm is 4  
(c) order cannot be found from the given information  
(d) None of these
- Q 4. A thin film with index of refraction 1.50 coats a glass lens with index of refraction 1.80. What is the minimum thickness of the thin film that will strongly reflect light with wavelength 600 nm ?  
(a) 150 nm                      (b) 200 nm  
(c) 300 nm                      (d) 450 nm
- Q 5. A thin film with index of refraction 1.33 coats a glass lens with index of refraction 1.50. Which of the following choices is the smallest film thicknesses that will not reflect light with wavelength 640 nm ?  
(a) 160 nm  
(b) 240 nm  
(c) 360 nm  
(d) 480 nm
- Q 6. A thick film of liquid polymer  $n = 1.25$  coats a slab of Pyrex,  $n = 1.50$ . White light is incident perpendicularly to the film. In the reflections, full destructive interference occurs for  $\lambda = 600$

nm and full constructive interference occurs for  $\lambda = 700$  nm. What is the thickness of the polymer film ?

- (a) 120 nm (b) 280 nm  
(c) 460 nm (d) 840 nm

### COMPREHENSION(Q.7 to Q.9)

Figure shows two flat glass plates  $P_1$  and  $P_2$  placed nearly (but not exactly) parallel forming an air wedge. The plates are illuminated normally by monochromatic light and viewed from above. Light waves reflected from the upper and lower surfaces of the air wedge give rise to an interference pattern.



- Q 7. Separation between successive bright fringes is  
(a)  $\frac{\lambda\ell}{4s}$  (b)  $\frac{\lambda\ell}{3s}$  (c)  $\frac{\lambda\ell}{s}$  (d)  $\frac{\lambda\ell}{2s}$
- Q 8. At line joining two plates, there is  
(a) maxima  
(b) minima  
(c) neither maxima nor minima  
(d) None of these
- Q 9. What will be the answer of last question if space between plates is filled by water  
(a) maxima  
(b) minima  
(c) neither maxima nor minima  
(d) None of these
- Q 10. A Fraunhofer's single slit diffraction is observed in the focal plane of a lens of focal length one metre. Slit width is 0.3 mm. The 2nd minima is observed at a distance of 4mm from the central maxima. Then the wave length of the light used is  
(a)  $5000\text{\AA}$  (b)  $6000\text{\AA}$   
(c)  $6500\text{\AA}$  (d)  $5400\text{\AA}$
- Q 11. Fraunhofer diffraction pattern of a single slit is obtained in the focal plane of lens of focal length 1m. If third maximum is formed at a distance of 5mm from the central maximum and wavelength of light used is  $5000\text{\AA}$ , then width of the slit will be –  
(a) 0.02 cm (b) 0.03 cm  
(c) 0.04 cm (d) 1 cm
- Q 12. If  $I_0$  is the intensity of the principle maximum in the single slit diffraction pattern, then what will be its intensity when the slit width is doubled ?  
(a)  $2I_0$  (b)  $4I_0$  (c)  $I_0$  (d)  $I_0/2$
- Q 13. A slit of width  $d$  is placed in front of a lens of focal length 0.5 m and is illuminated normally with light of wavelength  $5.89 \times 10^{-7}$  m. The first diffraction minima on either side of the central diffraction maximum are separated by  $2 \times 10^{-3}$  m. The width  $d$  of the slit is



- (a)  $2.94 \times 10^{-4} \text{m}$
- (b)  $4.94 \times 10^{-4} \text{m}$
- (c)  $5.94 \times 10^{-4} \text{m}$
- (d)  $6.94 \times 10^{-4} \text{m}$

PRATEEK JAIN  
PHYSICSAHOLICS

## Answer Key

Q.1 d	Q.2 a	Q.3 a,b,c	Q.4 b	Q.5 c
Q.6 d	Q.7 d	Q.8 b	Q.9 b	Q.10 b
Q.11 c	Q.12 b	Q.13 a		