

DPP – 7 (Electrostatics)

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/51>

Video Solution on YouTube:-


<https://youtu.be/zJhd11QYsfo>

Written Solution on Website:-


<https://physicsaholics.com/note/notesDetailis/40>

- Q 1. In a region with a uniform electric field, the number of lines of force per unit area is E . If a spherical metallic conductor is placed in the area, the field inside the conductor will be :
- (a) zero (b) E
 (c) more than E (d) less than E

- Q 2. A metallic solid sphere is placed in a uniform electric field. The lines of force follow the path(s) shown in figure as:




- (a) 1 (b) 2 (c) 3 (d) 4
- Q 3. A small conducting spherical shell with inner radius a and outer radius b is concentric with a larger conducting spherical shell with inner radius c and outer radius d (as shown in Fig). The inner shell has total charge $+2q$ and the outer shell has charge $+4q$. Calculate the electric field in terms of q and the distance r from the common centre of the two shells for: $a < r < b$.




- (a) zero (b) $\frac{1}{4\pi\epsilon_0} \frac{2q}{r^2}$
 (c) $-\frac{1}{4\pi\epsilon_0} \frac{6q}{r^2}$ (d) $\frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$
- Q 4. A solid conducting sphere of radius a has a net positive charge $2Q$. A conducting spherical shell of inner radius b and outer radius c is concentric with the solid sphere

and has a net charge $-Q$. The surface charge density on the inner and outer surfaces of the spherical shell will be:




- (a) $-\frac{2Q}{4\pi b^2}, \frac{Q}{4\pi c^2}$ (b) $-\frac{Q}{4\pi b^2}, \frac{Q}{4\pi c^2}$
 (c) zero, $\frac{Q}{4\pi c^2}$ (d) none of the above

Q 5. Figure shows three concentric conducting spherical shells with inner and outer shells earthed and the middle shell is given a charge q . The final charge on shell 1 and 3 are:



- (a) $\frac{4}{25}q, \frac{21}{25}q$ (b) $\frac{4}{25}q, -\frac{21}{25}q$
 (c) $-\frac{4}{25}q, -\frac{21}{25}q$ (d) none of the above

Q 6. Three concentric conducting spherical shells of radii $R, 2R$ and $3R$ carry charges $Q, -2Q$ and $3Q$, respectively. Find the electric potential at $r = R$:



- (a) $\frac{1}{4\pi\epsilon_0} \frac{Q}{R}$ (b) $-\frac{1}{4\pi\epsilon_0} \frac{Q}{R}$
 (c) $\frac{1}{4\pi\epsilon_0} \frac{3Q}{R}$ (d) none of the above




- Q 7. Three concentric conducting spherical shells of radii R , $2R$ and $3R$ carry charges Q , $-2Q$ and $3Q$, respectively. Compute the magnitude of electric field at $r = \frac{5}{2}R$: (where r is the radial distance from the centre)

Q) Three concentric conducting spherical shells of radii R , $2R$ and $3R$ carry charges Q , $-2Q$ and $3Q$, respectively. Compute the magnitude of electric field at $r = \frac{5}{2}R$: (where r is the radial distance from the centre)

- (a) $\frac{Q}{4\pi\epsilon_0 R^2}$ (b) $\frac{Q}{24\pi\epsilon_0 R^2}$
 (c) $\frac{Q}{25\pi\epsilon_0 R^2}$ (d) $\frac{Q}{45\pi\epsilon_0 R^2}$


- Q 8. Three concentric conducting spherical shells carry charges as $+4Q$ on the inner shell, $-2Q$ on the middle shell and $-5Q$ on the outer shell. The charge on the inner surface of the outer shell is:
- (a) Q (b) $4Q$
 (c) $-Q$ (d) $-2Q$

- Q 9. Find charge on outer surface of spherical shell-2 after joining the inner most shell and outer most shell by a conducting wire:




- (a) $\frac{3Q}{2}$ (b) $-\frac{3Q}{2}$
 (c) $-2Q$ (d) Q

- Q 10. Two conducting hollow spherical shells of radii R and $2R$ carry charges $-Q$ and $3Q$ respectively. How much charge will flow into the earth if inner shell is grounded ?




- (a) $\frac{3Q}{2}$ (b) $-\frac{3Q}{2}$
 (c) $\frac{Q}{2}$ (d) $-\frac{Q}{2}$

- Q 11. Figure shows three large metallic plates with charges $-Q$, $3Q$ and Q respectively. Determine the final charge on face C:



- (a) $\frac{5Q}{2}$ (b) $-\frac{5Q}{2}$
 (c) $\frac{Q}{2}$ (d) $-\frac{Q}{2}$

- Q 12. Two large, parallel conducting plates X and Y, kept close to each other, are given charges Q_1 and Q_2 ($Q_1 > Q_2$). The four surfaces of the plates are A, B, C and D, as shown in figure. Then:



- (a) The charge on A is $\frac{1}{2}(Q_1 + Q_2)$ (b) The charge on B is $\frac{1}{2}(Q_1 - Q_2)$
 (c) The charge on C is $-\frac{1}{2}(Q_1 - Q_2)$ (d) All of the above are correct



- Q 13. How does the charge densities of conductors vary on an irregularly shaped conductor?
- (a) Less at sharp and high at flat portion
 - (b) High at sharp and less at flat portion
 - (c) Remains constant
 - (d) Zero at sharp and high at flat portion

Answer Key

Q.1 a	Q.2 d	Q.3 a	Q.4 a	Q.5 c
Q.6 a	Q.7 c	Q.8 d	Q.9 b	Q.10 c
Q.11 a	Q.12 d	Q.13 b		