



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/99>

Video Solution on YouTube:-

https://youtu.be/_1dp4g7M5Dk

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/54>

- Q 1. Two particles of masses m and $2m$ are at a distance $3r$ apart at the ends of a straight-line AB . C is the center of mass of the system. The magnitude of the gravitational intensity due to the masses at C is
- (a) zero (b) $\frac{7Gm}{4r^2}$
(c) $\frac{9Gm}{4r^2}$ (d) $\frac{3Gm}{2r^2}$
- Q 2. The distance of the centers of moon and the earth is D . The mass of the earth is 81 times the mass of the moon. At what distance from the center of the earth, the gravitational force will be zero:
- (a) $\frac{D}{2}$ (b) $\frac{2D}{3}$
(c) $\frac{4D}{3}$ (d) $\frac{9D}{10}$
- Q 3. A point mass M is at a distance S from an infinitely long and thin rod of linear density D . If G is the gravitational constant, then gravitational force between the point mass and the rod is
- (a) $2 \frac{MGD}{S}$ (b) $\frac{MGD}{S}$
(c) $\frac{MGD}{2S}$ (d) $\frac{2MGD}{3S}$
- Q 4. The gravitational field due to a solid sphere of radius R and mass M at a point distant $R/2$ from the center of the sphere is
- (a) zero (b) $\frac{GM}{2R^2}$
(c) $\frac{GM}{R^2}$ (d) $\frac{4GM}{R^2}$
- Q 5. The height above the surface of earth at which the gravitational field intensity is reduced to 1% of its value on the surface of earth is: [R_e = radius of earth]
- (a) $100R_e$ (b) $10R_e$
(c) $99R_e$ (d) $9R_e$
- Q 6. The mass of the moon is 734×10^{20} kg and the radius is 1.74×10^6 m. The gravitational field strength at its surface is :
- (a) 1.45 N/kg (b) 1.55 N/kg
(c) 1.75 N/kg (d) 1.62 N/kg




Q 7. If earth is supposed to be a sphere of radius R , if g_{30} is value of acceleration due to gravity at latitude of 30° and g at the equator, the value of $g - g_{30}$ [ω = angular velocity of rotation of earth about its axis, R = radius of earth]

- (a) $\frac{5}{4}\omega^2 R$ (b) $\frac{3}{4}\omega^2 R$
 (c) $\omega^2 R$ (d) $\frac{1}{4}\omega^2 R$

Q 8. A tunnel is dug along a diameter of the earth. The force on a particle of mass m placed in the tunnel at a distance x from the center is: [M_e = mass of earth, R = radius of earth]



- (a) $\frac{GM_em}{R^3} x$ (b) $\frac{GM_em}{R^2} x$
 (c) $\frac{GM_em}{R^3 x}$ (d) $\frac{GM_e}{R^3 x}$

Q 9. A uniform metal sphere of radius a and mass M is surrounded by a thin uniform spherical shell of equal mass and radius $4a$ (figure) The center of the shell falls on the surface of the inner sphere. P_1 is at a distance $4a$ from center of metal sphere. Find the gravitational field at the point P_2 shown in the figure




- (a) $\frac{61GM}{900a^2}$ (b) $\frac{16GM}{3a^2}$
 (c) $\frac{35GM}{161a^2}$ (d) $\frac{51GM}{90a^2}$



Q 10. Two identical spherical balls each of mass m are placed as shown in figure. Plot the variation of g (gravitational intensity) along the x -axis.




(a)



(b)



- Q 11. At what height from surface of earth the gravitational field reduces by 75 % the gravitational field at the surface of earth ?
 (a) R (b) 2R
 (c) 3R (d) 4R
- Q 12. The gravitational field in a region is given by $\vec{E} = 5\hat{i} + 12\hat{j}$ (in N/Kg). Find the magnitude of the gravitational force acting on a particle of mass 2kg placed at the origin.
 (a) 26 N (b) 30 N
 (c) 20 N (d) 35 N
- Q 13. The gravitational field in a region is $(10\hat{i} - 10\hat{j})$ N/kg. Find the work done by gravitational force to shift slowly a particle of mass 1kg from point (1m, 1m) to a point (2m, -2m).
 (a) 10 J (b) -10 J
 (c) -40 J (d) 40 J
- Q 14. Two planets have the same average density, but their radii are R_1 and R_2 . If acceleration due to gravity on these planets be g_1 and g_2 respectively, then
 (a) $\frac{g_1}{g_2} = \frac{R_1}{R_2}$ (b) $\frac{g_1}{g_2} = \frac{R_2}{R_1}$
 (c) $\frac{g_1}{g_2} = \frac{R_1^2}{R_2^2}$ (d) $\frac{g_1}{g_2} = \frac{R_1^3}{R_2^3}$
- Q 15. Let the acceleration due to gravity be g_1 at a height h above the earth's surface, and g_2 at a depth d below the earth's surface. If $g_1 = g_2$, $h \ll R$ and $d \ll R$ then
 (a) $h = d$ (b) $h = 2d$
 (c) $2h = d$ (d) it is not possible for g_1 to be equal to g_2
- Q 16. What would be the value of acceleration due to gravity at a point 5 km below the earth's surface?
 ($R_e = 6400\text{km}$, $g_E = 9.8 \text{ m/s}^2$)
 (a) 9.6 m/s^2 (b) 9.79 m/s^2
 (c) 9.89 m/s^2 (d) 10 m/s^2
- Q 17. What will be the acceleration due to gravity at a distance of 3200 km below the surface of the earth ? [Take $R_e = 6400 \text{ km}$]
 (a) 2.7 m/s^2 (b) 4.9 m/s^2
 (c) 9.8 m/s^2 (d) 19.6 m/s^2
- Q 18. From a solid sphere of mass M and radius R, a solid sphere of radius R/2 is removed as shown. Find gravitational force on mass m as shown



- (a) $\frac{5}{12} \frac{GMm}{R^2}$ towards left
- (b) $\frac{7}{36} \frac{GMm}{R^2}$ towards left
- (c) $\frac{3}{17} \frac{GMm}{R^2}$ towards right
- (d) $\frac{9}{11} \frac{GMm}{R^2}$ towards right

PRATEEK JAIN
PHYSICSAHOLICS

Answer Key

Q.1 b	Q.2 d	Q.3 a	Q.4 b	Q.5 d
Q.6 d	Q.7 d	Q.8 a	Q.9 a	Q.10 a
Q.11 a	Q.12 a	Q.13 d	Q.14 a	Q.15 c
Q.16 b	Q.17 b	Q.18 b		