



## DPP – 1 & 2 (Gravitation)

Video Solution on Website :-

<https://physicsaholics.com/home/courseDetails/100>

Video Solution on YouTube:-

<https://youtu.be/iqllbHGVv0I>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/55>

Q 1. Three particles of equal mass  $m$  are situated at the vertices of an equilateral triangle of side  $l$ . What should be the velocity of each particle, so that they move on a circular path without changing  $l$  -

(a)  $\sqrt{\frac{GM}{2l}}$

(b)  $\sqrt{\frac{GM}{l}}$

(c)  $\sqrt{\frac{2GM}{l}}$

(d)  $\sqrt{\frac{GM}{3l}}$

Q 2. A spherical shell is cut into two pieces along a chord as shown in figure . If  $I_1$  and  $I_2$  are gravitational field strength at P due to upper part and lower part respectively, then

(a)  $I_1 > I_2$

(b)  $I_1 < I_2$

(c)  $I_1 = I_2 = 0$

(d)  $I_1 = I_2 \neq 0$


Q 3. The figure represents a solid uniform sphere of mass  $M$  and radius  $R$ . A spherical cavity of radius  $r$  is at a distance  $a$  from the center of the sphere. The gravitational field inside the cavity is

(a) non - uniform

(b) towards the center of the cavity

(c) directly proportional to  $a$

(d) All of these



Q 4. Inside a uniform sphere of density  $\rho$  there is a spherical cavity whose center is at a distance  $l$  from the center of the sphere. Find the strength of the gravitational field inside the cavity.

(a)  $E = -\frac{2}{3}\pi G\rho l$

(b)  $E = -\frac{4}{3}\pi G\rho l$

(c)  $E = -\frac{4}{3}\pi^2 G\rho l$

(d)  $E = -\frac{4}{3}\pi G\rho^2 l^2$

Q 5. A straight rod of length  $l$  extends from  $x = a$  to  $x = L + a$ . Find the gravitational force exerts on a point mass  $m$  at  $x = 0$  is (if the linear density of rod  $\mu = A + Bx^2$ )

(a)  $Gm \left[ \frac{A}{a} + BL \right]$

(b)  $Gm \left[ A \left( \frac{1}{a} - \frac{1}{a+L} \right) + BL \right]$

(c)  $Gm \left[ BL + \frac{A}{a+L} \right]$


(d)  $Gm \left[ BL - \frac{A}{a} \right]$



- Q 6. The gravitational field in a region is given by  $(2\hat{i} + 2\hat{j})$  N/kg. What is the work done by an external agent to slowly shift a particle of mass 10 kg from the point (0,0) to a point (5m, 4m) ?
- (a) 180 J (b) - 180 J  
(c) 90 J (d) - 90 J


- Q 7. A small body of superdense material, whose mass twice the mass of the earth but whose size is very small compared to the size of the earth, starts from rest at a height  $H \ll R$  above the earth's surface, and reaches the earth's surface in time  $t$ . Then  $t$  is equal to-
- (a)  $\sqrt{2H/g}$  (b)  $\sqrt{H/g}$   
(c)  $\sqrt{2H/3g}$  (d)  $\sqrt{4H/3g}$


- Q 8. Two concentric spherical shells have masses  $m_1$  and  $m_2$  and radii  $r_1$  and  $r_2$ . Then-



- (a) Outer shell will have no contribution in gravitational field at point P  
(b) Force on P is directed towards O  
(c) Force on P is  $\frac{Gm_1m_2}{r^2}$   
(d) Force on P is  $\frac{Gm_1m_3}{r^2}$
- Q 9. A particle of mass  $m$  is placed at centre of uniform ring of mass  $M$  and radius  $R$ . Mass  $m$  is slightly displaced along axis and released. If ring is also free to move, angular frequency of shm is
- (a)  $\sqrt{\frac{G(M+m)}{R^3}}$  (b)  $\sqrt{\frac{G(M+m)}{2R^3}}$   
(c)  $\sqrt{\frac{GM(M+m)}{mR^3}}$  (d)  $\sqrt{\frac{Gm(M+m)}{MR^3}}$
- Q 10. Gravitational field at surface of earth is  $9.8\text{m/sec}^2$  and at height  $h$  it is  $9.6\text{m/sec}^2$ . Gravitational field at depth  $h$  from ground is
- (a)  $9.6\text{m/sec}^2$   
(b)  $9.7\text{m/sec}^2$   
(c)  $9.4\text{m/sec}^2$   
(d)  $10\text{m/sec}^2$

- Q 11. A sphere of mass  $M$  and radius  $R_2$  has a concentric cavity of radius  $R_1$  as shown in figure. The force  $F$  exerted by the sphere on a particle of mass  $m$  located at a distance  $r$  from the centre of sphere varies as  $(0 \leq r \leq \infty)$  :





- Q 12. The Earth may be regarded as a spherically shaped uniform core of density  $\rho_1$  and radius  $R/2$  surrounded by a uniform shell of thickness  $R/2$  and density  $\rho_2$ . Find the ratio of  $\frac{\rho_1}{\rho_2}$  if the value of acceleration due to gravity is the same at surface as at depth  $R/2$  from the surface
- (a) 2/1  
 (b) 5/3  
 (c) 7/4  
 (d) 7/3
- Q 13. A small body of mass  $m$  is projected with a velocity just sufficient to make it reach from the surface of a planet (of radius  $2R$  and mass  $3M$ ) to the surface of another planet (of radius  $R$  and mass  $M$ ). The distance between the centers of the two spherical planets is  $6R$ . Find distance of small body from centre of bigger planet when it acquires its minimum speed
- (a)  $2R[3 - \sqrt{3}]$   
 (b)  $3R[2 - \sqrt{3}]$   
 (c)  $2R[2 - \sqrt{3}]$   
 (d)  $3R[3 - \sqrt{3}]$
- Q 14. There is a smooth tunnel along a chord of earth. Mass of earth is  $M$  and its radius is  $R$ . Length of tunnel is  $R/2$ . A particle is released in tunnel from surface of earth (one end of tunnel). Velocity of particle at centre of tunnel is (assuming particle is just fitted in tunnel)
- (a)  $\frac{1}{2} \sqrt{\frac{GM}{R}}$                       (b)  $\frac{1}{4} \sqrt{\frac{GM}{R}}$   
 (c)  $\frac{1}{3} \sqrt{\frac{GM}{R}}$                       (d)  $\frac{1}{5} \sqrt{\frac{GM}{R}}$
- Q 15. A uniform spherical shell is divided into two hemispheres as shown in figure. P is a point at dividing surface (not at centre of sphere). Gravitational field at P due to lower hemisphere have direction along
- (a) a  
 (b) b  
 (c) c  
 (d) d



PRATEEK JAIN  
PHYSICSAHOLICS

## Answer Key

Q.1 b	Q.2 d	Q.3 c	Q.4 b	Q.5 b
Q.6 b	Q.7 c	Q.8 a, b, d	Q.9 a	Q.10 b
Q.11 b	Q.12 d	Q.13 d	Q.14 b	Q.15 c