



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/74>

Video Solution on YouTube:-

<https://youtu.be/BBdrwh3cjko>


Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/56>

- Q 1. Earth's magnetic field always has a horizontal component except at or Horizontal component of earth's magnetic field remains zero at
- (a) Equator (b) Magnetic poles  
(c) A latitude of  $60^\circ$  (d) An altitude of  $60^\circ$
- Q 2. The correct relation is
- (a)  $B = \frac{B_H}{B_V}$  (b)  $B = B_H \times B_V$   
(c)  $B = \sqrt{B_H^2 + B_V^2}$  (d)  $B = B_H + B_V$   
(Where  $B_H$  = Horizontal component of earth's magnetic field,  $B_V$  = Vertical component of earth's magnetic field and B = Total intensity of earth's magnetic field)
- Q 3. At a certain place, the horizontal component of earth's magnetic field is  $\frac{1}{\sqrt{3}}$  times of its vertical component. The angle of dip at that place is
- (a)  $30^\circ$  (b)  $45^\circ$   
(c)  $75^\circ$  (d)  $60^\circ$
- Q 4. The earth's magnetic field at a certain place has a horizontal component 0.3 Gauss and the total strength 0.5 Gauss. The angle of dip is:
- (a)  $\tan^{-1} \frac{3}{4}$  (b)  $\sin^{-1} \frac{3}{4}$   
(c)  $\tan^{-1} \frac{4}{3}$  (d)  $\sin^{-1} \frac{3}{5}$
- Q 5. A magnetic needle (free to rotate in any direction) will show which one of the following directions at the earth's magnetic pole
- (a) Vertical  
(b) No particular direction  
(c) Bent at  $45^\circ$  to the vertical  
(d) Horizontal
- Q 6. A short magnet of moment  $6.75 \text{ Am}^2$  produces a neutral point on its axis. If horizontal component of earth's magnetic field is  $5 \times 10^{-5} \text{ Wb/m}^2$ , then the distance of the neutral point from center of magnet should be
- (a) 10 cm (b) 20 cm  
(c) 30 cm (d) 40 cm



- Q 7. At a given place on earth's surface the horizontal component of earth's magnetic field is  $2 \times 10^{-5} \text{T}$  and resultant magnetic field is  $4 \times 10^{-5} \text{T}$ . The angles of dip at this place is:
- (a)  $30^\circ$  (b)  $60^\circ$   
(c)  $90^\circ$  (d)  $45^\circ$
- Q 8. The true value of angle of dip at a place is  $60^\circ$ , the apparent dip in a plane inclined at an angle of  $30^\circ$  with magnetic meridian is
- (a)  $\tan^{-1}\left(\frac{1}{2}\right)$  (b)  $\tan^{-1}(2)$   
(c)  $\tan^{-1}\left(\frac{2}{3}\right)$  (d) None of these
- Q 9. The true dip at a place is  $30^\circ$ . What is the apparent dip when the dip circle is turned  $60^\circ$  out of the magnetic meridian ?
- (a)  $\tan^{-1}\left(\frac{1}{\sqrt{6}}\right)$  (b)  $\tan^{-1}\left(\frac{\sqrt{3}}{2}\right)$   
(c)  $\tan^{-1}\left(\frac{2}{\sqrt{3}}\right)$  (d) None of these
- Q 10. If a dip circle is placed in a vertical plane at an angle of  $30^\circ$  to the magnetic meridian, the dip needle makes an angle of  $45^\circ$  with the horizontal. The real dip at that place is?
- (a)  $\tan^{-1}\left(\frac{2}{3}\right)$  (b)  $\tan^{-1}\left(\frac{\sqrt{3}}{2}\right)$   
(c)  $\tan^{-1}\left(\frac{2}{\sqrt{3}}\right)$  (d)  $\tan^{-1}\left(\frac{\sqrt{3}}{\sqrt{2}}\right)$
- Q 11. At a certain place a magnet makes 30 oscillations per minute. At another place where the magnetic field is double, its time period will be
- (a) 4 sec (b) 2 sec  
(c)  $\frac{1}{2}$  sec (d)  $\sqrt{2}$  sec
- Q 12. If the strength of the magnetic field is increased by 21%, the frequency of a magnetic needle oscillating in that field:
- (a) Increases by 10% (b) Decreases by 10%  
(c) Increases by 11% (d) Decreases by 11%
- Q 13. A magnetic needle of magnetic moment  $6.7 \times 10^{-2} \text{Am}^2$  and moment of inertia  $7.5 \times 10^{-6} \text{kg m}^2$  is performing simple harmonic oscillations in a magnetic field of 0.01T. Time taken for 10 complete oscillations is :
- (a) 7.76 s (b) 6.65 s  
(c) 8.89 s (d) 9.98 s
- Q 14. Time period for a magnet is T. If it is divided in four equal parts along its axis and perpendicular to its axis as shown then time period for each part will be



- (a)  $4T$  (b)  $\frac{T}{4}$   
 (c)  $\frac{T}{2}$  (d)  $T$

- Q 15. Two bar magnets of the same mass, length and breadth but magnetic moments  $M$  and  $2M$  respectively, when jointed in same position (Similar pole in same direction), time period is 3 sec. What will be the time period when they are placed in different position (Similar pole in opposite direction) :
- (a)  $\sqrt{3}$  sec (b)  $3\sqrt{3}$  sec  
 (c) 3 sec (d) 6 sec
- Q 16. A dip needle in a vertical plane perpendicular to the magnetic meridian will remain-
- (a) Vertical  
 (b) Horizontal  
 (c) In any direction  
 (d) Inclined at  $45^\circ$  with horizontal
- Q 17. When a magnet is placed vertical then the number of neutral point obtained in the plane of paper is-
- (a) 1 (b) 2 (c) 4 (d) 3
- Q 18. The magnetic needle of a tangent galvanometer is deflected at an angle  $30^\circ$ . The horizontal component of earth's magnetic field  $0.34 \times 10^{-4}$  T is along the plane of the coil. The magnetic field of coil-
- (a)  $1.96 \times 10^{-4}$  T (b)  $1.96 \times 10^{-5}$  T  
 (c)  $1.96 \times 10^4$  T (d)  $1.96 \times 10^5$  T

## Answer Key

|        |        |        |        |        |
|--------|--------|--------|--------|--------|
| Q.1 b  | Q.2 c  | Q.3 d  | Q.4 c  | Q.5 a  |
| Q.6 c  | Q.7 b  | Q.8 b  | Q.9 c  | Q.10 b |
| Q.11 d | Q.12 a | Q.13 b | Q.14 c | Q.15 b |
| Q.16 a | Q.17 a | Q.18 b |        |        |