



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/99>

Video Solution on YouTube:-

<https://youtu.be/k9B1ZnbEv-E>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/54>

- Q 1. An astronaut orbiting the earth in a circular orbit 120 km above the surface of earth, gently drops a spoon out of space-ship. The spoon will
- (a) Fall vertically down to the earth
  - (b) Move towards the moon
  - (c) Will move along with space-ship
  - (d) Will move in an irregular way then fall down to earth
- Q 2. A satellite of mass  $m$ , initially at rest on the earth, is launched into a circular orbit at a height equal to the radius of the earth. The minimum energy required is
- (a)  $\frac{\sqrt{3}}{4}mgR$
  - (b)  $\frac{1}{2}mgR$
  - (c)  $\frac{1}{4}mgR$
  - (d)  $\frac{3}{4}mgR$
- Q 3. A period of geostationary satellite is
- (a) 24 hr
  - (b) 12 hr
  - (c) 30 hr
  - (d) 48 hr
- Q 4. A satellite of mass  $m$  is orbiting close to the surface of the earth Radius  $R=6400$  km has kinetic energy  $K$ . The corresponding kinetic energy of the satellite to escape from the earth's gravitational field is
- (a)  $K$
  - (b)  $2K$
  - (c)  $mgR$
  - (d)  $mK$
- Q 5. Two satellites of masses  $m_1$  and  $m_2$  ( $m_1 > m_2$ ) are revolving the earth in circular orbits of radius  $r_1$  and  $r_2$  ( $r_1 > r_2$ ) respectively. Which of the following statement is true regarding their velocities  $V_1$  and  $V_2$
- (a)  $V_1 = V_2$
  - (b)  $V_1 < V_2$
  - (c)  $V_1 > V_2$
  - (d)  $\frac{V_1}{r_1} = \frac{V_2}{r_2}$
- Q 6. A geostationary satellite can be installed :
- (a) Over the north or south pole
  - (b) Over any city on the equator
  - (c) In an orbit marking an angle of  $43.3^\circ$  with the equator plane
  - (d) In an orbit marking an angle of  $66.5^\circ$  with the equatorial plane



- Q 7. An artificial satellite revolves round the earth at a height of 1000km. The radius of the earth is  $6.38 \times 10^3$  km. Mass of the earth =  $6 \times 10^{24}$ kg,  $G = 6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$ . Find the orbital speed and period of revolution of the satellite
- (a) 7364 m/s, 6297 s                      (b) 1064 m/s, 6297 s  
(c) 9364 m/s, 9297 s                      (d) 8364 m/s, 7297 s
- Q 8. A satellite S is moving in an elliptical orbit around the earth. The mass of the satellite is very small compared to the mass of the earth. Then
- (a) the acceleration of `S` is always directed towards the center of the earth  
(b) the angular momentum of `S` about the center of the earth changes in direction, but its magnitude remains constant  
(c) the total mechanical energy of `S` varies periodically with time  
(d) the linear momentum of `S` remains constant in magnitude
- Q 9. If an artificial satellite is moving in a circular orbit around earth with speed equal to one fourth of  $V_e$  from earth, then height of the satellite above the surface of the earth is ( $R$  = radius of earth)
- (a) 7R    (b) 4R  
(c) 3R    (d) R
- Q 10. An artificial satellite is moving in a circular orbit around the earth with a speed equal to half the magnitude of escape velocity from the earth. If the satellite is stopped suddenly in its orbit and allowed to fall freely onto the earth, find the speed with which it hits the surface of the earth. ( $R$  = radius of earth)
- (a)  $\sqrt{gR}$     (b)  $\sqrt{\frac{gR}{2}}$   
(c)  $\sqrt{\frac{g}{R}}$     (d)  $\sqrt{\frac{gM}{R}}$
- Q 11. The orbital speed of an artificial satellite very close to the surface of the earth is  $V_0$ . Then the orbital speed of another artificial satellite at a height equal to three times the radius of the earth from surface is
- (a)  $4V_0$     (b)  $2V_0$   
(c)  $0.5V_0$     (d)  $4V_0$
- Q 12. A satellite with kinetic energy  $E_k$  is revolving round the earth in a circular orbit. How much more kinetic energy should be given to it so that it may just escape into outer space
- (a)  $E_k$     (b)  $2E_k$   
(c)  $\frac{1}{2}E_k$     (d)  $3E_k$
- Q 13. If  $V_e$  and  $V_0$  represent the escape velocity and orbital velocity of a satellite corresponding to a circular orbit of radius R, then
- (a)  $V_e = V_0$     (b)  $\sqrt{2}V_0 = V_e$   
(c)  $V_e = \frac{V_0}{\sqrt{2}}$     (d)  $V_e$  and  $V_0$  are not related



Q 14. The angular velocity of rotation of a star (mass  $M$  and radius  $R$ ), such that the matter will start escaping from its equator is:

- (a)  $\sqrt{\frac{2GR}{M}}$                       (b)  $\sqrt{\frac{GM}{R^3}}$   
(c)  $\sqrt{\frac{2GM}{R}}$                       (d)  $\sqrt{\frac{2GM^2}{R}}$

Q 15. The escape velocity on a planet is equal to the orbital velocity (on surface) on another planet. If the radii of two planets are equal, determine the ratio of acceleration due to gravity on the two planets

- (a) 5 : 2                              (b) 3 : 4  
(c) 2 : 3                              (d) 1 : 2

PRATEEK JAIN  
PHYSICSAHOLICS

## Answer Key

Q.1 c	Q.2 d	Q.3 a	Q.4 b	Q.5 b
Q.6 b	Q.7 a	Q.8 a	Q.9 a	Q.10 a
Q.11 c	Q.12 a	Q.13 b	Q.14 b	Q.15 d