




<b>DPP-3</b> (Gravitation)						
Video Solution on Website :-		https://physicsaholics.com/home/courseDetails/100				
Video Solution on YouTube:-		https://youtu.be/u6xHsalyQkA				
Written Solution on Website:-		https://physicsaholics.com/note/notesDetalis/55				
Q 1. Q 2.	increase in potential en (a) mgR (b) $\frac{1}{2}$ m A small block of superc towards earth. Find its	hgR (c) 2 mgR (d) $\frac{1}{4}$ mgR dense material has mass 2 × 10 <sup>24</sup> kg. It is at a height h << R. It falls s speed when it is at a height h/2 -				
Q 3.	(a) $\sqrt{\frac{2gh}{3}}$ (b) $\sqrt{\frac{3gh}{4}}$ (c) $\sqrt{\frac{3gh}{5}}$ (d) $\sqrt{\frac{gh}{2}}$ Which of the following statement(s) is/are correct (a) An astronaut going from Earth to Moon will experience weightlessness once (b) When a thin uniform spherical shell gradually shrinks maintaining its shape, the gravitational potential at the centre decreases (c) In the case of spherical shell, the plot of potential versus distance from centre is continuous					
Q 4. (	<ul> <li>(d) In the case of spherical shell, the plot of gravitational field intensity I versus distance from centre is continuous</li> <li>Q 4. A person brings a mass of 1kg from infinity to a point A. Initially the mass was at rest but it moves at a speed of 2m/s as it reaches A. The work done by the person on the mass is -3J. The potential at A is- <ul> <li>(a) -3J/kg</li> <li>(b) -2 J/kg</li> <li>(c) -5 J/kg</li> <li>(d) none of these</li> </ul> </li> </ul>					

Q 5. A point P lies on the axis of a ring of mass M and radius a, at a distance a from its centre C. A small particle starts from P and reaches C under gravitational attraction only. Its speed at C will be -

(a) 
$$\sqrt{\frac{2GM}{a}}$$
  
(b)  $\sqrt{\frac{2GM}{a}\left(1-\frac{1}{\sqrt{2}}\right)}$   
(c)  $\sqrt{\frac{2GM}{a}\left(\sqrt{2}-1\right)}$   
(d) zero

Q 6. A body is projected vertically upwards from the earth's surface to reach a height 7R, where R is the Radius of earth. The velocity required to do so is-

(a) 
$$\sqrt{\frac{7GM}{8R}}$$
 (b)  $\sqrt{\frac{7GM}{4R}}$  (c)  $\sqrt{\frac{8GM}{3R}}$  (d)  $\sqrt{\frac{20}{11}\frac{GM}{R}}$ 





Q 7. A hole is drilled from the surface of earth to its centre. A particle is dropped from rest at the surface of earth. The speed of the particle when it reaches the centre of the earth in terms of its escape velocity on the surface of earth ve is: (b)  $v_e$ 

(a) 
$$\frac{v_e}{2}$$

(c)  $\sqrt{2}v_{e}$ 


- (d)  $\frac{v_e}{\sqrt{2}}$
- Q 8. The potential at the surface of a planet of mass M and radius R is assumed to be zero. Choose the most appropriate option:
  - (a) The potential at infinity is  $\frac{GM}{R}$
  - (b) The potential at the centre of planet is  $-\frac{GM}{R}$
  - (c) Both (a) and (b) are correct
  - (d) Both (a) and (b) are wrong
- Q 9. Two bodies of masses  $m_1$  and  $m_2$  are initially at rest placed infinite distance apart. They are then allowed to move towards each other under mutual gravitational attraction. Their relative velocity when they are r distance apart is :

(a) 
$$\sqrt{\frac{2G(m_1+m_2)}{r}}$$
 (b)  $\sqrt{\frac{2Gm_1m_2}{(m_1+m_2)r}}$ 

(c)

- $Q\,10$ . A particle is projected from surface of a planet with speed v in vertically upward direction . This particle reaches to infinity with zero speed. If we project same particle at angle  $\pi/4$  with vertical with same speed v
  - (a) It will move on elliptical path
  - (b) It will move to infinity with zero velocity
  - (c) It will move to infinity with nonzero velocity
  - (d) It will move on parabolic path
- A fixed gaseous planet attracts a light body from infinity. Velocity of light body at infinite is Q 11. zero. Body reaches to surface of planet with speed v. if resistive forces are negligible , velocity of body at centre of planet will be
  - (a) v
  - (b) 2v
  - (c) 3v/2 (d)  $\int \frac{3v}{2}$
- A tunnel is dug along a chord of Earth having length  $\sqrt{3}R$  is radius of Earth. A small block is Q 12. released in the tunnel from the surface of Earth. The particle comes to rest at the center of tunnel. Find coefficient of friction between the block and the surface of tunnel. Ignore the effect of rotation of Earth

(a) 0.22	(b) 0.67
(c) 0.86	(d) 0.99





## PRATIER JAMA PRATIER JAMAS PHYSICSALL PHYSICS

## **Answer Key**

Q.1 b	Q.2 b	Q.3 a, b, c	Q.4 c	Q.5 b
Q.6 b	Q.7 d	Q.8 c	Q.9 a	Q.10 b, d
Q.11 d	Q.12 c			