



## **DPP – 7 (Geometrical Optics)**

Video Solution on Website:-

https://physicsaholics.com/home/courseDetails/67


Video Solution on YouTube:-

https://youtu.be/iDVILtfceXw

Written Solution on Website:-

https://physicsaholics.com/note/notesDetalis/68


Select correct Statement for an object on principal axis of given arrangement Q 1.



- (a) If light is incident on surface 1 from left, the image formed after the first refraction is definitely virtual for a real object
- (b) If object is real and placed at left of surface 1, then the final image formed after two refractions may be real
- (c) If object is real and placed at left of surface 1, then the final image formed after two refractions may be virtual
- (d) If light is incident on surface 1 from left, the image formed after the first refraction is definitely real for a real object.

## **COMPREHENSION (Q2 to Q3)**

An extended object of size 2 cm is placed at a distance of d (cm) in medium (refractive index n = 3) from pole, on the principal axis of a spherical curved surface. The medium on the other side of refracting surface is air (refractive index n = 1).




- Q 2. For d = 20 cm, the distance of the image from the pole is
  - (a) 2 cm
- (b) 3 cm
- (d) 5 cm

- For d = 20 cm, the size of image is (a)  $\frac{1}{6}$  cm (b)  $\frac{2}{15}$  cm Q 3.
- $(c) \frac{6}{5} cm$
- $(d) \frac{3}{2} cm$
- Q 4. A plastic hemisphere has a radius of curvature of 8 cm and an index of refraction of 1.6. On the axis halfway between the plane surface and the spherical one (4 cm from each) is a small object O. The distance between the two images when viewed along the axis from the two sides of the hemisphere is approximately:




## hysicsaholics





- (a) 1.0 cm
- (b) 1.5 cm
- (c) 3.75 cm
- (d) 2.5 cm
- Q 5. A spherical surface of radius of curvature R, separates air (refractive index 1.0) from glass (refractive index 1.5). The center of curvature is in the glass. A point object P placed in air is found to have a real image Q in the glass. The line PQ cuts the surface at a point O and PO = OQ. The distance PO is equal to:
  - (a) 5 R
- (b) 3 R
- (c) 2 R
- (d) 1.5 R
- A point object is placed at the center of a glass sphere of radius 6 cm and refractive Q 6. index 1.5. The distance of the virtual image from the surface of the sphere is:
  - (a) 2 cm
- (b) 4 cm
- (c) 6 cm
- (d) 12 cm
- Parallel rays of light are falling on convex spherical surface of radius of curvature R = Q 7. 20 cm as shown. Refractive index of the medium is  $\mu = 1.5$ . After refraction from the spherical surface parallel rays:



- (a) actually, meet at some point
- (b) appears to meet after extending the refracted rays backwards
- (c) meet (or appears to meet) at a distance of 30 cm from the spherical surface
- (d) meet (or appears to meet) at a distance of 60 cm from the spherical surface
- For a spherical surface of radius of curvature R, separating two media of refractive Q 8. indices  $\mu_1$  and  $\mu_2$ , the two principal focal lengths are  $f_1$  and  $f_2$  respectively. Which one of the following relations is correct?

(a)  $f_1 = f_2$ (c)  $f_2 / \mu_2 = -f_1 / \mu_1$ 


- (b)  $f_2 / \mu_2 = f_1 / \mu_1$ (d)  $f_2 / \mu_1 = f_1 / \mu_2$
- Q9. A small object is enclosed in a sphere of solid glass 8 cm in radius. It is situated 2 cm from centre and is viewed from the side to which it is nearest. How far will it appear from the surface? ( $\mu_g = 3/2$ )
  - (a) 6 cm
- (b) 4 cm
- (c)  $5\frac{1}{3}$  cm
- (d)  $3\frac{2}{3}$  cm
- Q 10. A hollow glass sphere has outer diameter 4R and inner diameter 2R. A point object on the inner surface is viewed along the diameter from the opposite side. Find the distance between the object and its image.



## hysicsaholics



- (a)  $\frac{R(2\mu-1)}{3\mu-2}$
- (b)  $\frac{R(\mu-1)}{3\mu-2}$
- (c)  $\frac{R(2\mu-1)}{3\mu-1}$
- (d)  $\frac{R(\mu-1)}{3\mu-1}$
- Q 11. A small object of height 0.5 cm is placed in front of a convex surface of glass ( $\mu = 1.5$ ) of radius of curvature 10 cm. Find the height of the image formed in glass.



- (a) 2 cm
- (b) 1 cm
- (c) 3 cm
- (d) 4 cm
- Q 12. A parallel beam of light travelling in water (refractive index = 4/3) is refracted by a spherical air bubble of radius 2cm situated in water. Assuming the light rays to be paraxial, the position of the image due to refraction at the first surface is
  - (a) 6cm from the first surface
  - (b) 12 cm from the first surface
  - (c) 3cm from the first surface
  - (d) 10 cm from the first surface



Q.1 a, b, c	Q.2 c	Q.3 c	Q.4 d	Q.5 a
Q.6 c	Q.7 a, d	Q.8 c	Q.9 с	Q.10 d
Q.11 b	Q.12 a			