



## DPP – 1 (Theometry & Calorimetry)

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/54>

Video Solution on YouTube:-

<https://youtu.be/ywVgjA2pU4c>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetailis/79>

- Q 1. The freezing point on a thermometer is marked as  $-20^\circ$  and the boiling point as  $130^\circ$ . A temperature of human body ( $34^\circ\text{C}$ ) on this thermometer will be read as:  
(a)  $31^\circ$  (b)  $51^\circ$   
(c)  $20^\circ$  (d) none of these
- Q 2. In a temperature scale called Z, the boiling point of water is 65Z and freezing point is  $-14\text{Z}$ . Then the temperature  $T = -98\text{Z}$  corresponds on the Fahrenheit scale to  
(a)  $-191\text{F}$  (b)  $-159\text{F}$   
(c)  $79\text{F}$  (d) none of these
- Q 3. If a thermometer reads freezing point of water as  $20^\circ\text{C}$  and boiling point  $150^\circ\text{C}$ . How much thermometer reads when the actual temperature is  $60^\circ\text{C}$ ?  
(a)  $98^\circ\text{C}$  (b)  $110^\circ\text{C}$   
(c)  $40^\circ\text{C}$  (d)  $60^\circ\text{C}$
- Q 4. A centigrade and a Fahrenheit thermometers are dipped in boiling water. The water temperature is lowered until the Fahrenheit thermometer reads  $140^\circ\text{C}$ . The fall in temperature registered by centigrade thermometer is  
(a)  $80^\circ\text{C}$  (b)  $40^\circ\text{C}$   
(c)  $50^\circ\text{C}$  (d)  $90^\circ\text{C}$
- Q 5. 100 gm of ice at  $0^\circ\text{C}$  is mixed with 100 g of water at  $100^\circ\text{C}$ . What will be the final temperature of the mixture?  
(A)  $10^\circ\text{C}$  (B)  $20^\circ\text{C}$  (C)  $30^\circ\text{C}$  (D)  $40^\circ\text{C}$
- Q 6. A lead bullet of 10g travelling at 300 m/s strikes against a block of wood and comes to rest. Assuming 50% of heat is absorbed by the bullet, the increase in its temperature is (specific heat of lead =  $150\text{ J/kg}\cdot\text{K}$ )  
(a)  $100^\circ\text{C}$  (b)  $125^\circ\text{C}$  (c)  $150^\circ\text{C}$  (d)  $200^\circ\text{C}$
- Q 7. Equal masses of three liquids A, B and C have temperatures  $10^\circ\text{C}$ ,  $25^\circ\text{C}$  and  $40^\circ\text{C}$  respectively. If A and B are mixed, the mixture has a temperature of  $15^\circ\text{C}$ . If B and C are mixed, the mixture has a temperature of  $30^\circ\text{C}$ . If A and C are mixed, the mixture will have a temperature of  
(a)  $16^\circ\text{C}$  (b)  $20^\circ\text{C}$  (c)  $25^\circ\text{C}$  (d)  $29^\circ\text{C}$



- Q 8. On increasing temperature of water from freezing point to boiling point its specific heat  
(a) remains constant (b) first increases then decreases  
(c) first decreases then increases (d) decreases throughout
- Q 9. Three different liquids with equal masses ( $m$ ), specific heat as  $s_A$ ,  $s_B$  and  $s_C$  & initial temperature as  $T_A$ ,  $T_B$  &  $T_C$  are kept closed in a isolated container, then -  
(a) final temperature of mixture will be  $\frac{1}{3}(T_A + T_B + T_C)$  if  $s_A = s_B = s_C$   
(b) heat given by liquid A to liquid B & C will be  $\frac{ms_A}{3}(2T_A - T_B - T_C)$  if  $s_A = s_B = s_C$   
(c) heat absorbed by liquid C will be  $\frac{ms_C}{s_A+s_B+s_C}[s_A(T_A-T_C) + s_B(T_B - T_C)]$   
(d) heat absorbed by liquid A is  $\frac{ms_A}{3}(T_B + T_C - 2T_A)$  if  $s_A = s_B = s_C$

## Answer Key

Q.1 a	Q.2 b	Q.3 a	Q.4 b	Q.5 a
Q.6 c	Q.7 a	Q.8 c	Q.9 a,b,c,d	


PLUS **ICONIC\*\***

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months	₹2,333/mo	>
No cost EMI	₹56,000	
18 months	₹2,625/mo	>
No cost EMI	₹47,250	
12 months	₹3,208/mo	>
No cost EMI	₹38,500	
6 months	₹4,667/mo	>
No cost EMI	₹28,000	

To be paid as a one-time payment

[View all plans](#)

 Add a referral code APPLY

# PHYSICSLIVE


PLUS **ICONIC\*\***

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months	₹2,100/mo	>
No cost EMI	+10% OFF ₹50,400	
18 months	₹2,363/mo	>
No cost EMI	+10% OFF ₹42,525	
12 months	₹2,888/mo	>
No cost EMI	+10% OFF ₹34,650	
6 months	₹4,200/mo	>
No cost EMI	+10% OFF ₹25,200	

To be paid as a one-time payment

[View all plans](#)

 Awesome! **PHYSICSLIVE** code applied ✗

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS.

# Written Solution

**DPP-1 Calorimetry & Thermometry**

**By Physicsaholics Team**

## Solution: 1

$$\frac{X - n_1}{n_2 - n_1} = \frac{C - 0}{100}$$

Here  $n_1$  is freezing point &  $n_2$  is boiling point in X thermometer

$$\Rightarrow \frac{X - (-20)}{130 - (-20)} = \frac{34}{100}$$

$$\Rightarrow \frac{X + 20}{150} = \frac{34}{100}$$

$$\Rightarrow X + 20 = 51$$

$$\Rightarrow X = 31$$

Solution: 2

$$\frac{Z - n_1}{n_2 - n_1} = \frac{F - 32}{180}$$

$$\Rightarrow \frac{-98 - (-14)}{65 - (-14)} = \frac{F - 32}{180}$$

$$\Rightarrow \frac{-84}{79} = \frac{F - 32}{180}$$

$$\Rightarrow F = -159$$

(B)

ANS : b

Solution: 3

$$\frac{X - n_1}{n_2 - n_1} = \frac{C - 0}{180}$$

$$\Rightarrow \frac{X - 20}{150 - 20} = \frac{60}{180}$$

$$\Rightarrow \frac{X - 20}{130} = \frac{6}{18}$$

$$\begin{aligned} \Rightarrow X &= 78 + 20 \\ &= 98 \end{aligned}$$

ANS : a

Solution: 4

$$\frac{C - 0}{100} = \frac{F - 32}{180}$$

$$\Rightarrow \frac{C}{100} = \frac{140 - 32}{180}$$

$$\Rightarrow C = 60$$

Initial temperature in centigrade = 100

fall in temperature in centigrade  
=  $100 - 60 = 40$

(b)

ANS : b



Solution: 5)

Let final temperature is  $T^{\circ}\text{C}$

where  $0 \leq T < 100$

Heat given by water = Heat taken by ice

$$\Rightarrow 100 \times 1 \times (100 - T) = 100 \times 80 + 100 \times 1 (T - 0)$$

$$100 - T = 80 + T$$

$$2T = 20$$

$$T = 10^{\circ}\text{C}$$

(A)

ANS : a

Solution: 6

$$\begin{aligned} \text{K.E. of bullet} &= \frac{1}{2} \times 0.01 \times (300)^2 \\ &= 450 \text{ J} \end{aligned}$$

$$\begin{aligned} \text{Heat absorbed by bullet} &= \frac{450}{2} \\ &= 225 \text{ J} \end{aligned}$$

$$\Delta Q = m s \Delta T$$

$$225 = 0.01 \times 150 \Delta T$$

$$\Delta T = \frac{2250}{15} = 150^\circ\text{C}$$

(c)

ANS : c

Solution: 7

Let  $\theta_1, \theta_2$  &  $\theta_3$  are specific heats of A, B & C.

When A & B are mixed

$$m \theta_1 (15 - 10) + m \theta_2 (15 - 25) = 0$$

$$\Rightarrow 5\theta_1 - 10\theta_2 = 0 \Rightarrow \theta_1 = 2\theta_2 \quad \text{---(i)}$$

When B & C are mixed

$$m \theta_2 (30 - 25) + m \theta_3 (30 - 40) = 0$$

$$\Rightarrow 5\theta_2 - 10\theta_3 = 0 \Rightarrow \theta_2 = 2\theta_3 \quad \text{---(ii)}$$

When A & C are mixed, final temp. is T.

$$m s_1 (T-10) + m s_3 (T-40) = 0$$

$$\Rightarrow \frac{s_1}{s_3} (T-10) + T-40 = 0$$

$$\Rightarrow \frac{s_1}{s_2} \times \frac{s_2}{s_3} (T-10) + T-40 = 0$$

$$\Rightarrow 4T-40 + T-40 = 0$$

$$\Rightarrow 5T = 80 \Rightarrow T = 16^\circ\text{C}$$

(A)

ANS : a

## Solution: 8

Specific Heat of water first decreases then increases on increasing its temperature from  $0^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ . It is minimum near  $38^{\circ}\text{C}$ .

(c)

ANS : c

Solution: 9

$$\text{If } \rho_A = \rho_B = \rho_C = \rho$$

$$m \rho (T - T_A) + m \rho (T - T_B) + m \rho (T - T_C) = 0$$

$$\Rightarrow 3T = T_A + T_B + T_C$$

$$\Rightarrow T = \frac{T_A + T_B + T_C}{3} \quad \text{(A) is correct}$$

Heat given by A

$$\Delta Q_A = m \rho_A (T_A - T)$$

$$= m \rho_A \left( T_A - \frac{T_A + T_B + T_C}{3} \right)$$

$$= \frac{m \rho_A}{3} (2T_A - T_B - T_C)$$

$$\text{If } \rho_A \neq \rho_B \neq \rho_C$$

$$m \rho_A (T - T_A) + m \rho_B (T - T_B) + m \rho_C (T - T_C) = 0$$

$$\Rightarrow T = \frac{\rho_A T_A + \rho_B T_B + \rho_C T_C}{\rho_A + \rho_B + \rho_C}$$

Heat absorbed by C

$$\Delta Q_c = m \rho_C (T - T_C)$$

$$= m \rho_C \left[ \frac{\rho_A T_A + \rho_B T_B + \rho_C T_C}{\rho_A + \rho_B + \rho_C} - T_C \right]$$

$$= m \rho_C \left[ \frac{\rho_A (T_A - T_C) + \rho_B (T_B - T_C)}{\rho_A + \rho_B + \rho_C} \right]$$

(c) is correct

ANS : c

Heat absorbed by A if  $s_A = s_B = s_C = s$

$$\Delta Q_A = m s_A (T - T_A)$$

$$= m s_A \left[ \frac{T_A + T_B + T_C}{3} - T_A \right]$$

$$= \frac{m s_A}{3} [T_B + T_C - 2T_A]$$

(D) is correct



For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/54>

Video Solution  
on YouTube:-

<https://youtu.be/ywVgjA2pU4c>

Written Solution  
on Website:-

<https://physicsaholics.com/note/notesDetails/79>

 **SUBSCRIBE**



[@Physicsaholics](#)

[@Physicsaholics\\_prateek](#)

[@NEET\\_Physics](#)  
[@IITJEE\\_Physics](#)

[physicsaholics.com](#)

[Unacademy](#)



**CLICK**

Chalo Niklo