



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/60>


Video Solution on YouTube:-

<https://youtu.be/jWXHJMdv-3k>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/78>

- Q 1. A monoatomic gas is taken along path AB as shown. Calculate change in internal energy of system ?




- (a) 279.8 J (b) 341 J
(c) 241 J (d) None of these
- Q 2. Internal energy of ideal diatomic gas at 300 K is 100 J. In this 100 J
(a) Potential energy = 0
(b) Rotational kinetic energy = 40 J
(c) Translational kinetic energy = 60 J
(d) Translational kinetic energy = 100 J
- Q 3. The average degrees of freedom per molecule for a gas is 6. The gas performs 25 J of work when it expands at constant pressure. The change in internal energy of gas is
(a) 75 J (b) 100 J
(c) 150 J (d) 125 J
- Q 4. One mole of an ideal gas whose pressure changes with volume as $P = \alpha V$, where α is a constant, is expanded so that its volume increases η times. Find change in internal energy in terms of initial volume V and degree of freedom f ?
(a) $f\alpha V^2(\eta^2 - 1)/8$ (b) $f\alpha V^2(\eta^2 - 1)/4$
(c) $f\alpha V^2(\eta^2 - 1)/2$ (d) None of these
- Q 5. 5 mole of O_2 is heated at constant volume from $10^\circ C$ to $20^\circ C$. What is the change in its internal energy?
(a) 250 cal (b) 200 cal
(c) 100 cal (d) 400 cal
- Q 6. The internal energy U of the air in an open room is
(a) Higher in day (b) Higher in night
(c) Equal in day and night (d) None of the above
- Q 7. For an ideal gas,




- (a) the change in internal energy in a constant-pressure process from temperature T_1 to T_2 is equal to $nC_v(T_2 - T_1)$
- (b) the change in internal energy of the gas and the work done by the gas are equal in magnitude in an isobaric process
- (c) the internal energy does not change in an isothermal process
- (d) Change in internal energy in isochoric process from temperature T_1 to T_2 is equal to $nC_v(T_2 - T_1)$.

- Q 8. Slope of internal energy vs temperature graph will be highest for
- (a) O_2
 - (b) H_2
 - (c) NH_3
 - (d) He
- Q 9. Volume of a gas is decreased to half of its initial volume. Magnitude of change in internal energy will be minimum in process
- (a) Isobaric
 - (b) Isothermal
 - (c) Process having equation $PV^{-1} = \text{constant}$
 - (d) Process having equation $PV^{-2} = \text{constant}$
- Q 10. Relation between U , P and V for ideal gas is $U = 2 + 2PV$ then gas is
- (a) Mono-atomic
 - (b) Diatomic
 - (c) Poly-atomic
 - (d) Mixture of mono and diatomic
- Q 11. Pressure versus density graph of an ideal gas is shown in figure:



- (a) during the process DA work done by the gas is positive
 - (b) during the process DA work done by the gas is negative
 - (c) during the process BC Internal energy of the gas is increasing
 - (d) none of the above
- Q 12. One mole of an ideal monatomic gas (initial temperature T_0) is made to go through the cycle abc a shown in the figure. If U denotes the internal energy, then choose the correct alternatives :



- (a) $U_c - U_a = 10.5 RT_0$
- (b) $U_b - U_a = 4.5 RT_0$



- (c) $U_c > U_b > U_a$
(d) $U_c - U_b = 6 RT_0$

Answer Key

Q.1 d	Q.2 a,b,c	Q.3 a	Q.4 c	Q.5 a
Q.6 c	Q.7 a,c,d	Q.8 c	Q.9 b	Q.10 d
Q.11 a	Q.12 a,b,c,d			

PRATEEK JAIN
PHYSICSAHOLICS