

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/76>


Video Solution on YouTube:-

https://youtu.be/_jv92zQJKg


Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/81>

- Q 1. A particle of mass $4m$ which is at rest explodes into four equal fragments. All 4 fragments scattered in the same horizontal plane. Three fragments are found to move with velocity V each as shown in the figure. The total energy released in the process of explosion is :



- (a) $mV^2(3 - \sqrt{2})$ (b) $2mV^2$ (c) $mV^2 \frac{(3-\sqrt{2})}{2}$ (d) $mV^2 \frac{(1+\sqrt{2})}{2}$
- Q 2. A bomb of mass 30 kg at rest explodes into two pieces of masses 18 kg and 12 kg . The velocity of 18 kg mass is 6 ms^{-1} . The kinetic energy of the other mass is :
- (a) 256 J (b) 486 J (c) 524 J (d) 324 J
- Q 3. Two blocks of mass 3 kg and 6 kg respectively are placed on a smooth horizontal surface. They are connected by a light spring of force constant $k = 200\text{ N/m}$. Initially the spring is unstretched. The indicated velocities are imparted to the blocks. The maximum extension of the spring will be:




- (a) 30 cm (b) 25 cm (c) 20 cm (d) 15 cm
- Q 4. Two blocks A and B of mass m and $2m$ are connected by a massless spring of force constant k . They are placed on a smooth horizontal plane. Spring is stretched by an amount x and then released. The relative velocity of the blocks when the spring comes to its natural length is



- (a) $\left(\sqrt{\frac{3k}{2m}}\right)x$ (b) $\left(\sqrt{\frac{2k}{3m}}\right)x$ (c) $\sqrt{\frac{2kx}{m}}$ (d) $\sqrt{\frac{3kx}{2m}}$




- Q 5. An isolated particle of mass m is moving in horizontal plane (x - y), along the x -axis, at a certain height above the ground. It suddenly explodes into two fragments of masses $m/4$ and $3m/4$. An Instant later, the smaller fragment is at $y = +15$ cm. The larger fragment at this instant is at:
- (a) $y = -5$ cm (b) $y = +20$ cm (c) $y = +5$ cm (d) $y = -20$ m
- Q 6. In the arrangement shown in figure match the following:




| Table-1 | | Table-2 | |
|---------|---|---------|-------------|
| (A) | Velocity of centre of mass | (P) | 2 SI unit |
| (B) | Velocity of combined mass when compression in the spring is maximum | (Q) | 1 SI unit |
| (C) | Maximum compression in the spring | (R) | 4 SI unit |
| (D) | Maximum potential energy stored in the spring | (S) | 0.5 SI unit |

- Q 7. Two blocks A (3kg) and B (6kg) are connected by a spring of stiffness 512 N/m and placed on a smooth horizontal surface. Initially the spring has its equilibrium length. Velocities 1.8 m/s and 2.2 m/s are imparted to A and B in opposite direction. The maximum extension in the spring will be –



- (a) 25 cm (b) 10 cm (c) 12 cm (d) 2.5 cm
- Q 8. Two blocks A and B of mass m and $2m$ respectively are connected by a massless spring of spring constant K . This system lies over a smooth horizontal surface. At $t = 0$ the block A has velocity u towards right as shown while the speed of block B is zero, and the length of spring is equal to its natural length at that instant. In each situation of column-I, certain statements are given and corresponding results are given in column-II, Match the statements in column-I to the corresponding results in column-II :





Column I

- (A) The velocity of block A
- (B) The velocity of block B
- (C) The kinetic energy of system of two blocks
- (D) The potential energy of spring

Column II


- (P) Can never be zero
- (Q) may be zero at certain instants of time
- (R) is minimum at maximum compression of spring
- (S) is maximum at maximum extension of spring

Q 9. Two masses m and $2m$ are attached to two ends of an ideal spring and the spring is in the compressed state. The energy of spring is 60 joule. If the spring is released, then-



- (a) the energy of both bodies will be same
 - (b) energy of smaller body will be 10J
 - (c) energy of smaller body will be 20J
 - (d) energy of smaller body will be 40 J
- Q 10. In last problem if initial compression was 12 cm, then maximum displacement of $2m$ is
- (a) 4 cm
 - (b) 8 cm
 - (c) 6 cm
 - (d) 2 cm

Q 11. Two masses m and $2m$ are attached to two ends of an ideal spring of stiffness K and the spring is in its natural length. At $t = 0$, a constant force F starts acting on $2m$ in rightward direction. Maximum elongation in spring is



- (a) $2F/K$
 - (b) F/K
 - (c) $F/2K$
 - (d) $2F/3K$
- Q 12. A gun of mass $2m$ when fitted (fixed) with ground fires bullet of mass m with velocity v . What will be velocity of same bullet if gun is free to move ?
- (a) v
 - (b) $\frac{v}{\sqrt{2}}$
 - (c) $v\sqrt{3/2}$
 - (d) $\frac{v}{\sqrt{3}}$



Answer Key

| | | | | |
|--|---------------|--|--------------|---------------|
| Q.1 a | Q.2 b | Q.3 a | Q.4 a | Q.5 a |
| Q.6 (A) Q, (b) Q, (C) Q, (D) p | Q.7 a | Q.8 $A \rightarrow P; B \rightarrow Q; C$ $\rightarrow P, R; D \rightarrow Q, S$ | Q.9 d | Q.10 b |
| Q.11 d | Q.12 c | | | |