



## DPP – 4 (Heat Transfer)

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/68>

Video Solution on YouTube:-

[https://youtu.be/x\\_1ZvgeTzDU](https://youtu.be/x_1ZvgeTzDU)

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/83>

### Passage (Q.1 to Q.4)

The body radiates energy or cools down to surrounding depends on the temperature of the surrounding. It says that the rate of heat loss to the surrounding at higher temperature is more than that of the body at lower temperature. To perform that we take a metal ball of mass 1 kg is heated by means of a 20 W heater in a room at 20°C. The temperature of the ball rises continuously but the rate of increase in temperature decreases continuously and finally it becomes zero, when the temperature of the ball reaches to 50°C. Corresponding to above observation answer the following questions

- Q 1. Find the rate of loss of heat to the surrounding when the ball is at 50°C.  
(A) 10 W (B) 15 W  
(C) 20 W (D) 25 W
- Q 2. Using Newtons law of cooling, find rate of heat loss to the surrounding when ball is at 30°C ?  
(A)  $\frac{10}{3} W$  (B)  $\frac{20}{3} W$   
(C)  $\frac{30}{3} W$  (D)  $\frac{40}{3} W$
- Q 3. Assume that the temperature of the ball rises uniformly from 20°C to 30°C in 5 minutes find the total loss of heat to the surrounding during this period –  
(A) 250 J (B) 500 J  
(C) 750 J (D) 1000 J
- Q 4. Find the specific heat capacity of the metal  
(A) 500 J/kg-K (B) 1000 J/kg-K  
(C) 1500 J/kg-K (D) 2000 J/kg-K
- Q 5. An object is cooled from 75°C to 65°C in 2 minutes in room at 30°C. The time taken to cool the same object from 55°C to 45°C in the same room is  
(a) 5 minute  
(b) 3 minute  
(c) 4 minute  
(d) 2 minute
- Q 6. A planet having surface temperature T has solar constant S. An angle  $\theta$  is subtended by the sun at planet then  
(a)  $S \propto T^2$   
(b)  $S \propto T$



- (c)  $S \propto \theta^0$   
(d)  $S \propto \theta^2$

Q 7. Four spheres A, B, C and D of different metals but of same radius are kept at same temperature. The ratio of their densities and specific heats are 2 : 3 : 5 : 1 and 3 : 6 : 2 : 4. Which sphere will show the fastest rate of cooling (initially):

- (a) A (b) B  
(c) C (d) D


Q 8. A body cools from 50°C to 40°C in 5 minutes. The surrounding temperature is 20°C. In what further time (in minutes) will it cool to 30°C?

- (a) 5 (b) 15/2  
(c) 25/3 (d) 10

Q 9. A planet is at an average distance  $d$  from the sun, and its average surface temperature is  $T$ . Assume that the planet receives energy only from the sun, and loses energy only through radiation from its surface. Neglect atmospheric effects. If  $T \propto d^{-n}$ , the value of  $n$  is


- (a) 2 (b) 1  
(c) 1/2 (d) 1/4



Q 10. A body cools in a surrounding which is at a constant temperature of  $\theta_0$ . Assume that it obeys Newton's law of cooling. Its temperature  $\theta$  is plotted against time  $t$ . Tangents are drawn to the curve at the points P( $\theta = \theta_1$ ) and Q( $\theta = \theta_2$ ). These tangents meet the time axis at angles of  $\phi_2$  and  $\phi_1$ , as shown.



- (a)  $\frac{\tan \phi_2}{\tan \phi_1} = \frac{\theta_1 - \theta_0}{\theta_2 - \theta_0}$  (b)  $\frac{\tan \phi_2}{\tan \phi_1} = \frac{\theta_2 - \theta_0}{\theta_1 - \theta_0}$   
(c)  $\frac{\tan \phi_1}{\tan \phi_2} = \frac{\theta_1}{\theta_2}$  (d)  $\frac{\tan \phi_1}{\tan \phi_2} = \frac{\theta_2}{\theta_1}$

Q 11. If the temperature of a body ( $\theta$ ) is slightly more than the temperature of the surrounding ( $\theta_0$ ), then the rate of cooling is correctly represented by –





- Q 12. Two identical spheres A and B are suspended in an air chamber which is maintained at a temperature of  $50^{\circ}\text{C}$ . Find the ratio of heat lost per sec from the surface of A to that of B, if A and B are at temperature  $60^{\circ}$  and  $55^{\circ}\text{C}$  respectively.
- Q 13. A hot body placed in air is cooled according to Newton's law of cooling, the rate of decrease of temperature being  $K$  times the temperature difference from the surroundings. Starting from  $t = 0$ , the time in which the body loses half the maximum heat is given by  $\frac{\ln x}{K}$ , where  $x$  is equal to
- Q 14. Three discs A, B and C having radii 2 m, 4 m and 6 m respectively are coated with carbon black on their outer surfaces. The wavelengths corresponding to maximum intensity are 300 nm, 400 nm and 500 nm respectively. The power radiated by them are  $Q_A$ ,  $Q_B$  and  $Q_C$  respectively
- (a)  $Q_A$  is maximum                      (b)  $Q_B$  is maximum  
(c)  $Q_C$  is maximum                      (d)  $Q_A = Q_B = Q_C$
- Q 15. Maximum spectral radiance of black body corresponds to wavelength  $\lambda$ . If temperature is now changed so that maximum spectral radiance now corresponds to  $\frac{3\lambda}{4}$ . Then
- (a) New temperature is  $\frac{4}{3}$  times the old temperature  
(b) New temperature is  $\frac{3}{4}$  times the old temperature  
(c) Power radiated by body changes by factor  $\frac{256}{81}$   
(d) Power radiated by body changes by factor  $\frac{81}{256}$
- Q 16. A black body is at temperature of 2880 K. The energy of radiation emitted by this object between wavelength  $4990 \text{ \AA}$  and  $5000 \text{ \AA}$  is  $U_1$ ; between  $9990 \text{ \AA}$  and  $10000 \text{ \AA}$  is  $U_2$  and between  $14990 \text{ \AA}$  and  $15000 \text{ \AA}$  is  $U_3$ . The Wein's constant is  $b = 2.88 \times 10^{-3} \text{ mK}$ , Then
- (a)  $U_2 > U_1$                                   (b)  $U_2 > U_3$   
(c)  $U_1 = U_3 < U_2$                           (d)  $U_1 < U_2 < U_3$
- Q 17. Explanations of phenomena's in column-II is explained by laws given in column-I.

	Column I		Column II
(A)	Why days are hot and nights are cold in deserts	(P)	Wein's displacement law
(B)	Why blackened platinum wire when heated gradually appears red and then blue	(q)	Planck's law



(C)	Variation in spectral intensity with temperature in distribution of energy in black body spectrum	(r)	Kirchhoff's law
(D)	Determination of some stars being hotter than others	(s)	Stefan's law

- Q 18. Two bodies A and B have thermal emissivity's of 0.01 and 0.81 respectively. The outer surface areas of the two bodies are the same. The two bodies radiate energy at the same rate. The wavelength  $\lambda_B$ , corresponding to the maximum spectral radiancy in the radiation from B, is shifted from the wavelength corresponding to the maximum spectral radiancy in the radiation from A by  $1.00 \mu\text{m}$ . If the temperature of A is 5802 K,
- (a) the temperature of B is 1934 K
  - (b)  $\lambda_B = 1.5 \mu\text{m}$
  - (c) the temperature of B is 11604 K
  - (d) the temperature of B is 2901 K

## Answer Key

Q.1 c	Q.2 b	Q.3 d	Q.4 a	Q.5 c
Q.6 d	Q.7 d	Q.8 c	Q.9 c	Q.10 b
Q.11 d	Q.12 2	Q.13 2	Q.14 b	Q.15 a, c
Q.16 a, b	Q.17 A(r), B(p), C(q), D(p)	Q.18 a, b		