



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/107>


Video Solution on YouTube:-

<https://youtu.be/xWx4T0wumvY>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetailis/88>

- Q 1. Sum of four nonzero vectors is zero , Then
- (a) they must be coplanar
 - (b) they may be coplanar
 - (c) Sum of magnitude of any two vectors must be equal to that of other two vectors.
 - (d) magnitude of sum of any two vectors must be equal to that of other two vectors
- Q 2. if \vec{A} has greater magnitude than \vec{B} Then maximum possible angle between sum of vectors with \vec{A} is
- (a) $\cos^{-1} \frac{B}{A}$
 - (b) $\cos^{-1} \frac{B}{\sqrt{A^2+B^2}}$
 - (c) $\sin^{-1} \frac{B}{A}$
 - (d) $\sin^{-1} \frac{B}{\sqrt{A^2+B^2}}$
- Q 3. If $\vec{A} \times \vec{B} = \vec{A} \times \vec{C}$ then (all three vectors are nonzero vectors)
- (a) B must be equal to C .
 - (b) \vec{B} must be equal to \vec{C}
 - (c) \vec{B} and \vec{C} must be coplanar
 - (d) \vec{B} and \vec{C} must be colinear.
- Q 4. if $A = 1$, $B = 2$, $C = 3$ and angle between \vec{A} and \vec{B} , \vec{B} and \vec{C} , \vec{C} and \vec{A} are 60° each. Magnitude of resultant of \vec{A} , \vec{B} and \vec{C} is
- (a) 3
 - (b) 4
 - (c) 5
 - (d) 6
- Q 5. If $\vec{A} \cdot \vec{B} = \vec{A} \cdot \vec{C}$ then (all three vectors are nonzero vectors)
- (a) B must be equal to C .
 - (b) \vec{B} must be equal to \vec{C}
 - (c) \vec{B} and \vec{C} must be coplanar
 - (d) None of these
- Q 6. In an octagon ABCDEFGH of equal side, what is the sum of \vec{AB} , \vec{AC} , \vec{AD} , \vec{AE} , \vec{AF} , \vec{AG} , and \vec{AH}



- (a) $6 \vec{AO}$
- (b) $3 \vec{AO}$
- (c) $4 \vec{AO}$
- (d) \vec{AO}

Q 7. A particle is revolving in a circular track passing through point (3,4,5) about axis of rotation $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$. Radius of circle is

- (a) 1
- (b) $\sqrt{\frac{22}{7}}$
- (c) $\sqrt{\frac{32}{7}}$
- (d) $\sqrt{\frac{42}{7}}$

Q 8. In an equilateral DABC, AL, BM and CN are medians. Forces along BC and BA represented by them will have a resultant represented by -

- (a) $2\mathbf{AL}$
- (b) $2\mathbf{BM}$
- (c) $2\mathbf{CN}$
- (d) \mathbf{AC}

Q 9. A particle is moving with uniform velocity 10 m/sec from point (2 m,5m,6m) to (3m,7m,8m) Velocity vector of particle is

- (a) $\frac{10}{3}(\hat{i} + 2\hat{j} + 2\hat{k})$ m/sec
- (b) $\frac{10}{3}(\hat{i} + \hat{j} + 2\hat{k})$ m/sec
- (c) $\frac{5}{3}(\hat{i} + 2\hat{j} + 2\hat{k})$ m/sec
- (d) $\frac{5}{6}(\hat{i} + 2\hat{j} + 2\hat{k})$ m/sec

Q 10. Vector $\hat{i} + 2\hat{j} + 2\hat{k}$ is resolved in two rectangular components. One component is along $\hat{i} + \hat{j} + \hat{k}$. Other component is

- (a) $\sqrt{\frac{5}{3}}$
- (b) $\sqrt{\frac{1}{3}}$
- (c) $\sqrt{\frac{2}{3}}$
- (d) $\sqrt{\frac{2}{5}}$



Answer Key

Q.1) B,D	Q.2) C	Q.3) C	Q.4) C	Q.5) D
Q.6) D	Q.7) A	Q.8) B	Q.9) A	Q.10) C

PRATEEK JAIN
PHYSICSAHOLICS