

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/42>


Video Solution on YouTube:-

<https://youtu.be/rTg2MIwPV6g>

Written Solution on Website:-


<https://physicsaholics.com/note/notesDetailis/36>

- Q 1. If the velocity of block B in the given arrangement is 300 mm/sec towards right. Find the velocity of A:




- (a) 100 mm/sec (b) 200 mm/sec
(c) 300 mm/sec (d) 400 mm/sec

- Q 2. Find the velocity of block B when ring A is moving downward with velocity v :




- (a) $v \sin \theta$ (b) $\frac{v}{2} \sin \theta$
(c) $v \cos \theta$ (d) $\frac{v}{2} \cos \theta$

- Q 3. If block A is moving horizontally with velocity V_A , then find the velocity of block B at the instant as shown in fig:.




- (a) $\frac{hV_A}{2\sqrt{x^2+h^2}}$ (b) $\frac{xV_A}{\sqrt{x^2+h^2}}$
 (c) $\frac{xV_A}{2\sqrt{x^2+h^2}}$ (d) $\frac{hV_A}{\sqrt{x^2+h^2}}$

- Q 4. A cart is being pulled up the incline, using a motor M and an ideal pulley and ideal rope arrangement as shown in figure. Then the speed of point ' P ' of the string with which it moves so that the car moves up the inclined plane with a constant speed of $V_{cart} = 2 \text{ m/s}$ is (Incline is at rest):




- (a) 12 m/s (b) 3 m/s
 (c) 5 m/s (d) 6 m/s

- Q 5. In Fig. a ball of mass m_1 and a block of mass m_2 are joined together with an inextensible string. The ball can slide on a smooth horizontal surface. If V_1 and V_2 are the respective speeds of the ball and the block, then determine the constraint relation between velocities of the two.




- (a) $V_2 = V_1 \cos \theta$ (b) $V_1 = V_2 \cos \theta$
 (c) $V_1 = V_2$ (d) $V_2 = V_1 \sin \theta$

Q 6. Find $V_B = ?$




- (a) 10 m/s (b) 8 m/s
(c) 14 m/s (d) 6 m/s

Q 7. Determine the speed with which block B rises in Fig. if the end of the cord at A is pulled down with a speed of 2 m/s.




- (a) 4 m/s (b) 3 m/s
(c) $\frac{3}{2}$ m/s (d) $\frac{1}{2}$ m/s

Q 8. Two rings each of mass $M = 100 \text{ gm}$ are constrained to move along a fixed horizontal rod. An ideal string is connected with rings and block of mass $M_0 = 200 \text{ gm}$ is connected to the mid point of string. At a certain moment the mass m is moving downward with velocity $\sqrt{3} \text{ m/s}$. Find the speed of ring of M at the moment:




- (a) 4 m/s (b) 3 m/s
(c) 2 m/s (d) 1 m/s

Q 9. In the given figure, find the speed of pulley P –




- (a) $\frac{V}{2}$ (b) $2V \cos \theta$
 (c) $\frac{2V}{\cos \theta}$ (d) $\frac{V}{2 \sin \theta}$

Q 10. Figure shows a rod of length l resting on a wall and the floor. Its lower end A is pulled towards left with a constant velocity u . As a result of this, end A starts moving down along the wall. Find the velocity of the other end B downward when the rod makes an angle θ with the horizontal:



- (a) $u \tan \theta$ (b) $u \cot \theta$
 (c) $u \sin \theta$ (d) $u \cos \theta$

Q 11. The velocities of A and B are marked in the figure. Find the velocity of block C (assume that the pulleys are ideal and string inextensible)



- (a) 2 m/s (b) 4 m/s
 (c) 5 m/s (d) $\sqrt{10}$ m/s



Answer Key

Q.1 b	Q.2 c	Q.3 c	Q.4 d	Q.5 a
Q.6 a	Q.7 d	Q.8 d	Q.9 c	Q.10 b
Q.11 c				

PRATEEK JAIN
PHYSICSAHOLICS

Plus leaderboard

Based on educator activity in last 30 days

- 1 Prateek Jain
11.4M mins
- 2 Ajay Mishra (Akm)
6.3M mins
- 3 Shubh Karan Choudhary (Skc)
5.9M mins
- 4 Dr Amit Gupta
5.5M mins
- 5 Ramesh Sharda
4.9M mins
- 6 Sandeep Nodiyal
4.8M mins
- 7 Shailendra Tanwar
3.6M mins
- 8 Vishal Vivek
2.7M mins
- 9 Garima Goel
2.7M mins
- 10 Saurabh Sharma
2.6M mins
- 11 Dr S K Singh
2.6M mins
- 12 Nishant Varshney



Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS and learn from India's Top Faculties.

NEET UG subscription

PLUS ICONIC*

- India's Best Educators
- Interactive Live Classes
- Structured Courses & PDFs
- Live Tests & Quizzes
- Personal Coach
- Study Planner

24 months	₹2,100/mo	>
No cost EMI	+10% OFF ₹50,400	
18 months	₹2,363/mo	>
No cost EMI	+10% OFF ₹42,525	
12 months	₹2,888/mo	>
No cost EMI	+10% OFF ₹34,650	
6 months	₹4,200/mo	>
No cost EMI	+10% OFF ₹25,200	

To be paid as a one-time payment

[View all plans](#)

Awesome! **PHYSICSLIVE** code applied


Written Solution

Physics DPP

DPP-2 NLM: Constraint Relation

By Physicsaholics Team

Solution: 1



$$3r_A = r$$

$$r_B = \frac{3}{2} r$$

$$r_B = \frac{3}{2} r_A$$

$$2r_B = 3r_A$$

$$2V_B = 3V_A$$


$$V_B = \frac{3}{2} V_A$$

$$V_A = \frac{2}{3} V_B$$

$$V_A = \frac{2}{3} (300)$$

$$V_A = 200 \text{ mm/sec}$$

OR



$$P_A + P_B = 0$$

$$(3r) V_A \cos 0^\circ + 2r V_B \cos(180^\circ) = 0$$

$$3V_A - 2V_B = 0$$


$$3V_A = 2V_B$$

$$V_A = \frac{2}{3} V_B = \frac{2}{3} \times 300$$

$$V_A = 200 \text{ mm/s}$$

Ans. b

Solution: 2



$$P_A + P_B = 0$$

$$T V_A \cos(180^\circ) + T V_B \cos 0^\circ = 0$$

$$-T V_A \cos 0 + T V_B = 0$$


$$V_B = V_A \cos 0$$

$$V_A = v$$

$$\therefore V_B = v \cos 0$$

Ans. c

Solution: 3



$$P_A + P_B = 0$$

$$T V_A \cos(180^\circ) + 2T V_B = 0$$

$$-T V_A \cos\theta + 2T V_B = 0$$


$$V_B = \frac{V_A \cos\theta}{2}$$

$$\cos\theta = \frac{x}{\sqrt{x^2 + h^2}}$$

$$\therefore V_B = \frac{x V_A}{2 \sqrt{x^2 + h^2}}$$

Ans. c

Solution: 4



$$P_p + P_c = 0$$

$$T v_p (\sin 180^\circ) + 3T v_c (\sin 90^\circ) = 0$$

$$-T v_p + 3T v_c = 0$$


$$v_p = 3 v_c$$

$$v_p = 3 \times 2$$

$$v_p = 6\text{ m/s}$$

Ans. d

Solution: 5



$$P_1 + P_2 = 0$$


$$T \cdot V_1 \cos(180^\circ - \theta) + T V_2 = 0$$

$$-T V_1 \cos \theta + T V_2 = 0$$

$$V_2 = V_1 \cos \theta$$

Ans. a

Solution: 6



$$P_A + P_B = 0$$

$$T v_A \cos(180^\circ) + T v_B \cos(37^\circ) = 0$$

$$-v_A + v_B \cos(37^\circ) = 0$$


$$v_B \left(\frac{4}{5}\right) = v_A$$

$$v_B = v_A \frac{5}{4} = 8 \times \frac{5}{4}$$

$$v_B = 10 \text{ m/s}$$

Ans. a

Solution: 7



$$P_A + P_B = 0$$

$$T \uparrow_A (\cos 180^\circ) + 4T \uparrow_B (\cos 180^\circ) = 0$$


$$-T(2) - 4T \uparrow_B = 0$$

$$v_B = -\frac{2}{4} = -\frac{1}{2} \text{ m/s}$$

$$\boxed{v_B = \frac{1}{2} \text{ m/s}} \text{ upward,}$$

Ans. d

Solution: 8



$$P_1 + P_2 + P_3 = 0$$

$$V_0(T \cos 60^\circ) + T(\cos 60^\circ) + TV \cos 30^\circ + TV \cos 30^\circ = 0$$

$$-2TV \cos 60^\circ + 2TV \cos 30^\circ = 0$$


$$V = \frac{+V_0 \cos 60^\circ}{\cos 30^\circ}$$

$$V = \frac{V_0}{\sqrt{3}} = \frac{\sqrt{3}}{\sqrt{3}}$$

$$V = 1 \text{ m/s}$$

Ans. d

Solution: 9



$$P_1 + P_2 + P_3 = 0$$

$$2T(V) \cos(180^\circ) + 2T(V) \cos(180^\circ) + 2T \cos \theta V_p = 0$$

$$-2TV \cos(180^\circ) + 2TV_p \cos \theta = 0$$


$$-2V + V_p \cos \theta = 0$$

$$V_p \cos \theta = 2V$$

$$\boxed{V_p = \frac{2V}{\cos \theta}}$$

Ans. c

Solution: 10



Since length of rod is constant

velocity of A along rod = velocity of B along rod

$$\Rightarrow V \cos \theta = V_B \sin \theta$$

$$\Rightarrow V_B = V \cot \theta$$

$$x^2 + y^2 = l^2$$
$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$x V_A + y (-V_B) = 0$$


$$(-V_B) = -\frac{x}{y} V_A$$

$$V_B = \frac{x}{y} V_A = \cot \theta V_A$$

$$V_B = V_A \cot \theta = V \cot \theta$$

Ans. b

Solution: 11



v_x of block will be equal to v_B (\because both are in contact and moving together in a direction)

$$\therefore v_x = 3 \text{ m/s}$$

Now for v_y .

$$P_A + P_B + (P)_y = 0$$

$$v_A(2T) \cos(180^\circ) + 2T v_B \cos 0^\circ + T v_y \cos 180^\circ = 0$$

$$-2v_A + 2v_B - v_y = 0$$

$$v_y = 2v_B - 2v_A$$

$$v_y = 2 \times 3 - 2(1)$$

$$v_y = 4 \text{ m/s}$$

$$v = \sqrt{v_x^2 + v_y^2}$$

$$v = \sqrt{3^2 + 4^2}$$

$$v = \sqrt{25} \text{ m/s}$$

$$v = 5 \text{ m/s}$$

Ans. c

For Video Solution of this DPP, Click on below link

Video Solution
on Website:-

<https://physicsaholics.com/home/courseDetails/42>

Video Solution
on YouTube:-

<https://youtu.be/rTg2MIwPV6g>

Written Solution
on Website:-

<https://physicsaholics.com/note/notesDetails/36>

Chalo Niklo