



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/57>

Video Solution on YouTube:-

<https://youtu.be/6oH-54BLk88>

Written Solution on Website:-




<https://physicsaholics.com/note/notesDetails/32>

- Q 1. Four molecules have speeds 2 km/sec, 3 km/sec, 4 km/sec and 5 km/sec. The root mean square speed of these molecules (in km/sec) is:
(a) $\sqrt{\frac{27}{2}}$ (b) $\sqrt{27}$ (c) 3.5 (d) $3\sqrt{3}$
- Q 2. At what temperature will the particles in a sample of helium gas have an rms speed of 1 km/s?
(a) 160°C (b) 222 K (c) 160 K (d) 222°C
- Q 3. The temperature of a gas is increased from 27°C to such an extent that its rms speed be double the speed at 27°C. The final temperature will be
(a) 927°C (b) 250°C (c) 600°C (d) 1200°C
- Q 4. At what temperature is the root mean square speed of an atom in an argon gas cylinder equal to the rms speed of a helium gas atom at -20°C ? (atomic mass of Ar = 39.9 u, and of He = 4.0 u)
(a) $2.52 \times 10^3\text{K}$ (b) $2.52 \times 10^3\text{K}$
(c) $25.2 \times 10^3\text{K}$ (d) 25.2×10^3
- Q 5. N (< 100) molecules of a gas have velocities 1, 2, 3,..... N km/s respectively. Then ratio of rms speed and average speed is:
(Given: The sum of squares of the first n natural numbers = $\frac{n(n+1)(2n+1)}{6}$)
(a) 1 (b) $\sqrt{\frac{(2N+1)(N+1)}{6N}}$
(c) $\sqrt{\frac{(2N+1)(N+1)}{6}}$ (d) $2\sqrt{\frac{(2N+1)}{6(N+1)}}$
- Q 6. Find the ratio of the mean speed of hydrogen molecules to the mean speed of nitrogen molecules in a sample containing a mixture of the two gases
(a) 14 (b) $\sqrt{14}$ (c) $\frac{1}{28}$ (d) $\frac{1}{\sqrt{14}}$
- Q 7. The mean speed of the molecules of a hydrogen sample equals the mean speed of the molecules of a helium sample. Calculate the ratio of the temperature of the hydrogen sample to the temperature of the helium sample
(a) $\frac{1}{2}$ (b) 2 (c) $\frac{1}{4}$ (d) 4



Q 8. The ratio of rms speed of an ideal gas molecules at pressure p to that at pressure $2p$ is
(a) $\frac{1}{2}$ (b) 2 (c) $\frac{1}{\sqrt{2}}$ (d) $\sqrt{2}$

Q 9. In a closed rigid container an ideal gas is filled. If the gas is heated, the graph of pressure (P) v/s root mean square speed (rms) will be :



(d) None of these

Q 10. A gas is filled in a rigid container at pressure P_0 . If the mass of each molecule is halved keeping the total number of molecules same and their r.m.s speed is doubled then find the new pressure

(a) $\sqrt{2}P_0$ (b) $3P_0$ (c) $\sqrt{3}P_0$ (d) $2P_0$

Q 11. At what temperature most probable speed of SO_2 molecule have the same value as root mean square speed of O_2 molecules at 300 K?

(a) 150K (b) 600K (c) 750K (d) 900K

Q 12. Most probable velocity, average velocity and root mean square velocity are related as:

(a) 1: 1.128: 1.224 (b) 1: 1.128: 1.424
(c) 1: 2.128: 1.224 (d) 1: 1.428: 1.442

Answer Key

Q.1 a	Q.2 c	Q.3 a	Q.4 b	Q.5 d
Q.6 b	Q.7 a	Q.8 c	Q.9 b	Q.10 d
Q.11 d	Q.12 a			


PLUS **ICONIC****

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months	₹2,333/mo	>
No cost EMI	₹56,000	
18 months	₹2,625/mo	>
No cost EMI	₹47,250	
12 months	₹3,208/mo	>
No cost EMI	₹38,500	
6 months	₹4,667/mo	>
No cost EMI	₹28,000	

To be paid as a one-time payment

[View all plans](#)

 Add a referral code APPLY

PHYSICSLIVE


PLUS **ICONIC****

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months	₹2,100/mo	>
No cost EMI	+10% OFF ₹50,400	
18 months	₹2,363/mo	>
No cost EMI	+10% OFF ₹42,525	
12 months	₹2,888/mo	>
No cost EMI	+10% OFF ₹34,650	
6 months	₹4,200/mo	>
No cost EMI	+10% OFF ₹25,200	

To be paid as a one-time payment

[View all plans](#)

 Awesome! **PHYSICSLIVE** code applied ✗

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS.

NEET & JEE Main Physics DPP- Solution

DPP- 2 Different type of Velocity and speed of gas molecules

By Physicsaholics Team

Solution 1:

$$V_{\text{rms}} = \sqrt{\frac{V_1^2 + V_2^2 + V_3^2 + \dots + V_n^2}{n}}$$

$$V_{\text{rms}} = \sqrt{\frac{(2)^2 + (3)^2 + (4)^2 + (5)^2}{4}}$$

$$V_{\text{rms}} = \sqrt{\frac{4 + 9 + 16 + 25}{4}}$$

$$V_{\text{rms}} = \sqrt{\frac{54}{4}}$$

$$V_{\text{rms}} = \sqrt{\frac{27}{2}} \text{ km/s}$$

Ans. a

Solution 2:

$$v_{\text{rms}} = \sqrt{\frac{3RT}{M}}$$

$$M_{\text{He}} = 4 \text{ gm or } 4 \times 10^{-3} \text{ kg}$$

$$(*) v_{\text{rms}} = 1 \text{ km/s} = 10^3 \text{ m/s}$$

$$(10^3)^2 = \frac{3 \times (8.31) \times T}{4 \times 10^{-3}}$$

$$10^6 = \frac{3 \times 8.31 \times T}{4 \times 10^{-3}}$$

$$T = \frac{10^6 \times 4 \times 10^{-3}}{3 \times 8.31} = 160 \text{ K}$$

$$\boxed{T \approx 160 \text{ K}}$$

Ans. c

Solution 3:

$$V_{\text{rms}} \propto \sqrt{T}$$

$$\frac{(V_{\text{rms}})_1}{(V_{\text{rms}})_2} = \sqrt{\frac{T_1}{T_2}}$$

$$\frac{v}{2v} = \sqrt{\frac{(273+27)}{T_2}}$$

$$\frac{1}{4} = \frac{300}{T_2}$$

$$T_2 = 1200 \text{ K}$$

⊙

$$T_2 = 927^\circ \text{C}$$

Ans. a

Solution 4:

$$V_r = \sqrt{\frac{3RT}{M}}$$

$$V_r \propto \sqrt{\frac{T}{M}}$$

$$\frac{V_{Ar}}{V_{He}} = \left(\sqrt{\frac{T}{M}}\right)_{Ar} \times \left(\sqrt{\frac{M}{T}}\right)_{He}$$

$\therefore V_{Ar} = V_{He}$ (given)

$$\frac{V_{Ar}}{V_{He}} = 1 = \left(\sqrt{\frac{T_{Ar}}{39.9}}\right) \times \left(\sqrt{\frac{4}{(273-20)}}\right)$$

$$(1)^2 = \frac{T_{Ar}}{39.9} \times \frac{4}{253}$$

$$T_{Ar} = 2523.077 \text{ K}$$

$$T_{Ar} = 2.52 \times 10^3 \text{ K}$$

Ans. b

Solution 5:

$$V_{avg} = \frac{1+2+3+4+\dots+N}{N}$$

$$\begin{aligned} S_n &= 1+2+3+\dots+N \\ &= \frac{N(N+1)}{2} \quad (\text{Sum of } N\text{-natural numbers}) \end{aligned}$$

$$\therefore V_{avg} = \frac{N(N+1)}{2N}$$

$$V_{avg} = \left(\frac{N+1}{2}\right) \text{ km/s}$$

Now

$$V_{rms} = \sqrt{\frac{1^2+2^2+3^2+4^2+\dots+N^2}{N}}$$

$$V_{rms} = \sqrt{\frac{N(N+1)(2N+1)}{6N}} \text{ km/s}$$

$$\frac{V_{rms}}{V_{avg}} = \frac{\sqrt{\frac{N(N+1)(2N+1)}{6N}}}{\left(\frac{N+1}{2}\right)}$$

$$= \frac{2}{(N+1)} \sqrt{\frac{(N+1)(2N+1)}{6}}$$

$$\frac{V_{rms}}{V_{avg}} = 2 \sqrt{\frac{(2N+1)}{6(N+1)}}$$

Ans. d

Solution 6:

$$v_m = \sqrt{\frac{8RT}{\pi M}}$$

$$v_m \propto \frac{1}{\sqrt{M}}$$

$$\frac{v_{H_2}}{v_{N_2}} = \sqrt{\frac{M_{N_2}}{M_{H_2}}} = \sqrt{\frac{28}{2}}$$

$$\frac{v_{H_2}}{v_{N_2}} = \sqrt{14}$$

Ans. b

Solution 7:

$$v_m = \sqrt{\frac{8RT}{\pi M}}$$

$$v_{H_2} = v_{He}$$
$$\sqrt{\frac{8RT_{H_2}}{M_{H_2}}} = \sqrt{\frac{8RT_{He}}{M_{He}}} \Rightarrow \left(\frac{8RT_{H_2}}{M_{H_2}} \right) = \left(\frac{8RT_{He}}{M_{He}} \right)$$

$$\frac{T_{H_2}}{T_{He}} = \frac{M_{He}}{M_{H_2}} = \frac{2}{4}$$

$$\frac{T_{H_2}}{T_{He}} = \frac{1}{2}$$

Ans. a

Solution 8:

$$V_{\text{rms}} = \sqrt{\frac{3P}{5}}$$

$$\frac{V_{\text{rms}}}{4} = \sqrt{\frac{P}{5}}$$

$$\frac{V_P}{V_{2P}} = \frac{\sqrt{P}}{\sqrt{2P}}$$

$$\frac{V_P}{V_{2P}} = \frac{1}{\sqrt{2}}$$

Ans. c

Solution 9:

closed container,

$$V = \text{constant}$$

$$\Rightarrow P \propto T$$


$$v_{rms} = \sqrt{\frac{3P}{\rho}}$$

$$v_{rms} \propto \sqrt{P}$$

$$v_{rms}^2 \propto P$$

$$\Rightarrow P \propto v_{rms}^2$$

↳ Parabolic curve between
P & v_{rms} . (upward open
parabola)



Ans. b

Solution 10:

$$v_{rms} = \sqrt{\frac{3P_0}{\rho}}$$

$$\rho = \frac{m}{V}$$

when mass of each molecule
is halved

then total mass = $m' = m/2$

$$\rho' = \frac{m'}{2V} = \frac{\rho}{2}$$

$$\text{So, } v'_{rms} = \sqrt{\frac{3P_1}{\rho'}}$$

$$\therefore v'_{rms} = 2 v_{rms} \quad (\text{given})$$

$$\sqrt{\frac{3P_1}{\rho'}} = 2 \sqrt{\frac{3P_0}{\rho}}$$

$$\frac{3P_1}{\rho'} = 4 \left(\frac{3P_0}{\rho} \right)$$

$$\frac{P_1}{\left(\frac{\rho}{2}\right)} = 4 \frac{P_0}{\rho}$$

$$2P_1 = 4P_0$$

$$\boxed{P_1 = 2P_0}$$

or

$$\boxed{P_1 = 2P_0}$$

Ans. d

Solution 11:

$$\text{Most Probable speed} = \sqrt{\frac{2RT}{M}}$$

$$\text{root Mean Square speed} = \sqrt{\frac{3RT}{M}}$$

$$\left(\sqrt{\frac{2RT}{M}}\right)_{\text{SO}_2} = \left(\sqrt{\frac{3RT}{M}}\right)_{\text{O}_2}$$

$$\sqrt{\frac{2R T_{\text{SO}_2}}{64}} = \sqrt{\frac{3R(300)}{32}}$$

$$\frac{2R T_{\text{SO}_2}}{64} = \frac{3R(300)}{32}$$

$$T_{\text{SO}_2} = 3 \times 300$$

$$T_{\text{SO}_2} = 900 \text{ K}$$

Ans. d

Solution 12:

$$\text{Most Probable speed} = v_m = \sqrt{\frac{2RT}{M}}$$

$$\text{Average velocity} = v_a = \sqrt{\frac{8RT}{\pi M}}$$

$$\text{RMS velocity} = v_{rms} = \sqrt{\frac{3RT}{M}}$$

$$v_m : v_a : v_{rms} = \sqrt{\frac{2RT}{M}} : \sqrt{\frac{8RT}{\pi M}} : \sqrt{\frac{3RT}{M}}$$

$$= \sqrt{2} : \sqrt{\frac{8}{\pi}} : \sqrt{3}$$

$$= 1 : \frac{1}{\sqrt{2}} \left(\sqrt{\frac{8}{\pi}} \right) : \frac{\sqrt{3}}{2}$$

$$= 1 : \sqrt{\frac{4}{\pi}} : \sqrt{\frac{3}{2}}$$

$$= 1 : 1.128 : 1.224$$

$$\boxed{v_m : v_a : v_{rms} = 1 : 1.128 : 1.224}$$

Ans. a

For Video Solution of this DPP, Click on below link

Video Solution
on Website:-

<https://physicsaholics.com/home/courseDetails/57>

Video Solution
on YouTube:-

<https://youtu.be/6oH-54BLk88>

Written Solution
on Website:-

<https://physicsaholics.com/note/notesDetails/32>

 **SUBSCRIBE**



[@Physicsaholics](#)

[@Physicsaholics_prateek](#)

[@NEET_Physics](#)
[@IITJEE_Physics](#)

[physicsaholics.com](#)

[Unacademy](#)



CLICK

Chalo Niklo