



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/47>

Video Solution on YouTube:-

<https://youtu.be/awMuuv1goWA>

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/48>

Q 1. Which of the following cylindrical rods, (given radius r and length l) each made of the same material and whose ends are maintained at the same temperature will conduct most heat?


(a) $r = 2r_0; l = 2l_0$

(b) $r = 2r_0; l = l_0$

(c) $r = r_0; l = 2l_0$

(d) $r = r_0; l = l_0$

Q 2. In the following situations, the length and area of cross-section of each rod is same. Find temperature θ at junction of rods




(a) $\frac{220}{3}^\circ\text{C}$

(b) $\frac{220}{5}^\circ\text{C}$

(c) $\frac{160}{5}^\circ\text{C}$

(d) $\frac{160}{3}^\circ\text{C}$

Q 3. Two identical square rods of metal are welded end to end as shown in figure (i), 20 calories of heat flows through it in 4 minutes. If the rods are welded as shown in figure (ii), the same amount of heat will flow through the rods in



(a) 1 minute

(b) 2 minutes

(c) 4 minutes

(d) 16 minutes

Q 4. The coefficient of thermal conductivity depends upon

(a) Temperature difference of two surfaces

(b) Area of the plate

(c) Thickness of the plate

(d) Material of the plate

Q 5. If the coefficient of conductivity of aluminium is $0.5 \text{ cal/cm-sec-}^\circ\text{C}$, then in order to conduct 10 cal/sec-cm^2 in steady state, the temperature gradient in aluminium must be:



- (a) $0.5\text{ }^{\circ}\text{C}/\text{cm}$ (b) $10\text{ }^{\circ}\text{C}/\text{cm}$
(c) $20\text{ }^{\circ}\text{C}/\text{cm}$ (d) $10.5\text{ }^{\circ}\text{C}/\text{cm}$
- Q 6. One end of a brass rod 2m long and having 1cm radius is maintained at $250\text{ }^{\circ}\text{C}$. When a steady state is reached, the rate of heat flow across any cross-section is 0.5 cal/s . What is the temperature of the other end ($K = 0.26\text{ cal/sec-cm-}^{\circ}\text{C}$)
(a) $100\text{ }^{\circ}\text{C}$ (b) $266.5\text{ }^{\circ}\text{C}$
(c) $127.5\text{ }^{\circ}\text{C}$ (d) 127.5 K
- Q 7. The length of a rod of aluminium is 1.0 m and its area of cross-section is 5.0 cm^2 . Its one end is kept at $250\text{ }^{\circ}\text{C}$ and the at $50\text{ }^{\circ}\text{C}$. How much heat will flow in the rod in 5.0 minutes . (Thermal conductivity 'K' for Al = $2.0 \times 10^{-1}\text{ KJs}^{-1}\text{m}^{-1}\text{ }^{\circ}\text{C}^{-1}$)
(a) 2000 J (b) 4000 J
(c) 6000 J (d) 8000 J
- Q 8. Find the thermal resistance of an aluminium rod of length 0.20 m and area of cross section $1 \times 10^{-4}\text{ m}^2$. The heat current is along the length of the rod. [Thermal conductivity of aluminium = 200 W/m-K]
(a) 10 k W^{-1} (b) 20 k W^{-1}
(c) 30 k W^{-1} (d) 40 k W^{-1}
- Q 9. Two rods A and B of same length and radius are joined together in series. the thermal conductivity of A and B are $2K$ and K . Under steady state conditions, if the temperature difference between the open ends of A and B is $36\text{ }^{\circ}\text{C}$, the temperature difference across 'A' is:
(a) $12\text{ }^{\circ}\text{C}$ (b) $18\text{ }^{\circ}\text{C}$
(c) $24\text{ }^{\circ}\text{C}$ (d) $9\text{ }^{\circ}\text{C}$
- Q 10. If the temperature difference between the two side of a wall is doubled, its thermal conductivity
(a) Is doubled (b) Is halved
(c) Become four times (d) None of these
- Q 11. A cotton sheet is ironed with hot electricity iron. How is energy transferred through the metal base of the iron to the sheet?
(a) By conduction (b) By convection only
(c) By radiation only (d) By convection & Radiation only
- Q 12. In a steady state the temperature of the ends A and B of a 20 cm long rod AB is $100\text{ }^{\circ}\text{C}$ and $0\text{ }^{\circ}\text{C}$. The temperature at the point C distant 9 cm from A is :
(a) $45\text{ }^{\circ}\text{C}$ (b) $55\text{ }^{\circ}\text{C}$
(c) $60\text{ }^{\circ}\text{C}$ (d) $65\text{ }^{\circ}\text{C}$



Answer Key

Q.1 b	Q.2 a	Q.3 a	Q.4 d	Q.5 c
Q.6 c	Q.7 c	Q.8 a	Q.9 a	Q.10 d
Q.11 a	Q.12 b			

PRATEEK JAIN
PHYSICSAHOLICS

Plus leaderboard

Based on educator activity in last 30 days

- 1 Prateek Jain
11.4M mins
- 2 Ajay Mishra (Akm)
6.3M mins
- 3 Shubh Karan Choudhary (Skc)
5.9M mins
- 4 Dr Amit Gupta
5.5M mins
- 5 Ramesh Sharda
4.9M mins
- 6 Sandeep Nodiyal
4.8M mins
- 7 Shailendra Tanwar
3.6M mins
- 8 Vishal Vivek
2.7M mins
- 9 Garima Goel
2.7M mins
- 10 Saurabh Sharma
2.6M mins
- 11 Dr S K Singh
2.6M mins
- 12 Nishant Varshney



Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS and learn from India's Top Faculties.

PLUS **ICONIC****

- India's Best Educators
- Interactive Live Classes
- Structured Courses & PDFs
- Live Tests & Quizzes
- Personal Coach
- Study Planner

24 months	₹2,100/mo	>
No cost EMI	+10% OFF ₹50,400	
18 months	₹2,363/mo	>
No cost EMI	+10% OFF ₹42,525	
12 months	₹2,888/mo	>
No cost EMI	+10% OFF ₹34,650	
6 months	₹4,200/mo	>
No cost EMI	+10% OFF ₹25,200	

To be paid as a one-time payment

[View all plans](#)


Awesome! **PHYSICSLIVE** code applied ✕

Written Solution

**DPP-1 Heat Transfer: Conduction: Rate of flow of heat,
Combination of rods (Series & Parallel)**

By Physicsaholics Team

Solution: 1



$$I_H = \frac{\kappa A \Delta \theta}{L} ; \text{ when } \Delta \theta = \text{constant}$$

$$\Rightarrow I_H \propto \frac{A}{L}$$

(i) $r = 2r_0$; $l = 2l_0$

$$I_{H1} \propto \frac{\pi (2r_0)^2}{2l_0} = \frac{2\pi r_0^2}{l_0}$$

(ii) $r = 2r_0$; $l = l_0$

$$I_{H2} \propto \frac{\pi (2r_0)^2}{l_0} = \frac{4\pi r_0^2}{l_0}$$

(iii) $r = r_0$; $l = 2l_0$

$$I_{H3} \propto \frac{\pi r_0^2}{2l_0}$$


(iv) $r = r_0$; $l = l_0$

$$I_{H4} = \frac{\pi r_0^2}{l_0}$$

$$\Rightarrow I_{H2} > I_{H1} > I_{H4} > I_{H3}$$

Ans. b

Solution: 2



$$\bar{I}_1 = \bar{I}_2$$

$$\frac{(2K)(A)(100 - \theta)}{L} = \frac{(K)(A)(\theta - 20)}{L}$$

$$2(100 - \theta) = (\theta - 20)$$



$$200 - 2\theta = \theta - 20$$

$$220^\circ\text{C} = 3\theta$$

$$\theta = \frac{220}{3}^\circ\text{C} \quad \text{Ans.}$$

Ans. a

Solution: 3



Case-1

$$I = \frac{\Delta\theta}{R}$$

$$R_{eq} = 2R$$

$$I_1 = \frac{\Delta\theta}{2R}$$

$$\frac{20}{4} = \frac{(100)}{2R} \quad \text{--- (1)}$$

Case-2

$$R_{eq} = R/2$$

$$I_2 = \frac{\Delta\theta}{R_{eq}} = \frac{100}{(R/2)}$$

$$\frac{20}{t} = \frac{100}{(R/2)} \quad \text{--- (2)}$$

$$\Rightarrow \frac{(1)}{(2)} \Rightarrow \frac{20/4}{20/t} = \frac{100/2R}{100/(R/2)}$$

$$\Rightarrow \frac{t}{4} = \frac{1}{4} \Rightarrow \boxed{t = 1 \text{ minute}} \text{ Ans.}$$

Ans. a

Solution: 4

Coefficient of Thermal Conductivity is the internal property of the metal and is independent of dimensions and other external factors. It only depend on nature of material.

PRATEEK JAIN
PHYSICSAHOLICS

Ans. d

Solution: 5

$$k = 0.5 \text{ cal/cm-sec-}^{\circ}\text{C}$$

$$I = \frac{kA(\Delta\theta)}{L}$$

$$\text{Temperature gradient} = \frac{\Delta\theta}{L}$$

$$\frac{\Delta\theta}{L} = \frac{I}{kA}$$

$$\text{given: } \frac{I}{A} = 10 \text{ cal/sec-cm}^2$$


$$\frac{\Delta\theta}{L} = \frac{I/A}{k} = \frac{10}{0.5}$$

$$\frac{\Delta\theta}{L} = \frac{100}{5} \Rightarrow$$

$$\frac{\Delta\theta}{L} = 20 \text{ }^{\circ}\text{C/cm} \text{ Ans.}$$

Ans. c

Solution: 6



$$I = 0.5 \text{ cal/s}$$

$$I = \frac{kA(250 - T)}{L}$$

$$0.5 = \frac{(0.26)(\pi \times (1)^2)(250 - T)}{200}$$


$$100 = (0.26)(\pi)(250 - T)$$

$$122.5 = 250 - T$$

$$\Rightarrow T = 127.5^\circ\text{C} \text{ Ans.}$$

Ans. c

Solution: 7

$$L = 1 \text{ m} = 100 \text{ cm}$$


$A = 5 \text{ cm}^2$

$2\pi \times 10^7 \text{ L}$ 50°C

$$t = 5 \text{ min.}$$

$$Q = ?$$

$$I = \frac{Q}{t} = \frac{k A (\Delta \theta)}{L}$$

$$\frac{Q}{(5 \times 60)} = \frac{(2 \times 10^7 \times 10^3) (5 \times 10^{-4}) (20)}{1}$$

$$Q = 5 \times 60 \times 2 \times 5 \times 20 \times 10^{-2}$$

$$Q = 5 \times 60 \times 2 \times 5 \times 2$$

$$Q = 6000 \text{ J}$$

Ans.

Ans. c

Solution: 8

$$I = \frac{kA \Delta \theta}{L} = \frac{\Delta \theta}{(L/kA)}$$

$$I = \frac{\Delta \theta}{R}$$

$$\Rightarrow R = \frac{L}{kA} = \frac{0.2}{200 \times 1 \times 10^{-4}}$$


$$R = \frac{2 \times 10^3}{2 \times 10^2}$$

$$R = 10 \text{ kW}^{-1}$$

Ans.

Ans. a

Solution: 9



$$I_A = I_B$$

$$\frac{(2k) A (\Delta\theta_A)}{L} = \frac{(k) A (\Delta\theta_B)}{L}$$

$$2 \Delta\theta_A = \Delta\theta_B$$

$$\Delta\theta_A + \Delta\theta_B = 36^\circ\text{C}$$

$$\Delta\theta_A + 2 \Delta\theta_A = 36^\circ\text{C}$$

$$3 \Delta\theta_A = 36^\circ\text{C}$$

$$\Delta\theta_A = 12^\circ\text{C} \quad \text{Ans.}$$

Ans. a

Solution: 10

Thermal conductivity is property of material it does not depend on temperature difference between two sides of wall. So, thermal conductivity will remains same.

(*It slightly varies with temperature but according to our syllabus we always consider constant with temperature change.)

PRATEEK JAIN
PHYSICSAHOLICS

Ans. d

Solution: 11

Conduction occurs when two objects at different temperatures are in contact with each other. Heat flows from the warmer to the cooler object until they are both at the same temperature. The iron and cotton sheet is in contact with each other so, there is conduction between them.

PRATEEK JAIN
PHYSICSAHOLICS

Ans. a

Solution: 12



$$I_{AC} = I_{CB}$$
$$\frac{KA(100-T)}{9 \text{ cm}} = \frac{KA(T-0)}{11 \text{ cm}}$$

$$11(100-T) = 9(T-0)$$

$$1100 - 11T = 9T$$

$$20T = 1100$$

$$T = \frac{110}{2}$$

$$T = 55^\circ\text{C} \quad \text{Ans.}$$

Temperature gradient
of rod = $G = \frac{100-0}{20}$

$$G = 5^\circ\text{C/cm.}$$

∴ Temp difference in length
AC

$$T_A - T_C = 5^\circ\text{C/cm} \times 9 \text{ cm}$$

$$100 - T_C = 45^\circ\text{C}$$

$$T_C = 55^\circ\text{C} \quad \text{Ans.}$$

Ans. b

For Video Solution of this DPP, Click on below link

Video Solution
on Website:-

<https://physicsaholics.com/home/courseDetails/47>

Video Solution
on YouTube:-

<https://youtu.be/awMuuv1goWA>

Written Solution
on Website:-

<https://physicsaholics.com/note/notesDetails/48>



[@Physicsaholics](#)

[@Physicsaholics_prateek](#)

[@NEET Physics](#)

[@IITJEE Physics](#)

[physicsaholics.com](#)



CLICK

Chalo Niklo