



Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/52>

Video Solution on YouTube:-

https://youtu.be/FOBVSJHhp_k

Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/74>


Q 1. On an inclined plane of inclination 30° , a ball is thrown at an angle of 60° with the horizontal from the foot of the incline with a velocity of $10\sqrt{3}$ m/s. If $g = 10$ m/s², then find the time in which ball will hit the inclined plane?

- (a) 1 s (b) 2 s (c) 3 s (d) 4 s

Q 2. Two bodies are projected from the same point with equal speeds in such a directions that they strike on the same point on a plane whose inclination is β . If α the angle of projection of the first, ratio of their times of flight is

- (a) $\frac{\sin \alpha}{\cos \beta}$ (b) $\frac{\sin \alpha}{\sin \beta}$ (c) $\frac{\sin(\alpha-\beta)}{\cos \beta}$ (d) $\frac{\sin(\alpha-\beta)}{\cos \alpha}$

Q 3. A baseball is projected with a velocity v making an angle θ with the incline of inclination α as shown in fig. Find the condition that the ball hits the incline at right angle.




- (a) $\cot \theta = 2 \tan \alpha$ (b) $\sin \theta = \cos \alpha$
(c) $\tan \theta = \sin \alpha$ (d) $\cot \theta = \cos \alpha$

Q 4. A projectile is required to hit a target whose coordinates relative to horizontal and vertical axes through the point of projection are (α, β) . If the gun velocity is $\sqrt{2g\alpha}$, it is impossible to hit the target if

- (a) $\beta > 3\alpha/4$ (b) $\beta \geq 1\alpha/4$
(c) $\beta \leq 3\alpha/4$ (d) $\beta \geq 3\alpha/4$

Q 5. A marble rolls down from top of a staircase with constant horizontal velocity 10 m/s. If each step is $y = 1$ meter high and $x = 1$ meter wide. To which step the marble will strike directly? ($g = 9.8$ m/s²)




- (a) 21st (b) 8th
 (c) 10th (d) 18th

Q 6. A particle is projected from origin of coordinate system. A target is fixed at point (40m, 30m). Find the minimum velocity of projectile to hit the target? ($g = 10 \text{ m/s}^2$)


- (a) 10 m/s (b) 17 m/s
 (c) $20\sqrt{2}$ m/s (d) $10\sqrt{5}$ m/s

Q 7. A particle is projected at an angle θ with an inclined plane making an angle β with the horizontal as shown in figure, speed of the particle is u , after time t find y component of velocity when particle is at maximum distance from the incline plane ?



- (a) $\frac{u}{\sqrt{2}}$ (b) $\frac{2u}{\sqrt{3}}$
 (c) $\frac{\sqrt{2}u}{3}$ (d) zero

Q 8. A ball is projected from point A with a velocity 10 m/s perpendicular to the inclined plane as shown in figure. Range of the ball on the inclined plane is :



- (a) $\frac{40}{3}$ m (b) $\frac{20}{13}$ m
 (c) $\frac{13}{20}$ m (d) $\frac{13}{40}$ m


Q 9. A plane surface is inclined making an angle θ with the horizontal. From the bottom of this inclined plane, a bullet is fired with velocity v . The maximum possible range of the bullet on the inclined plane is

- (a) $\frac{v^2}{g}$ (b) $\frac{v^2}{g(1+\sin \theta)}$
 (c) $\frac{v^2}{g(1-\sin \theta)}$ (d) $\frac{v^2}{g(1+\cos \theta)}$

Q 10. A ball is projected horizontal with a speed v from the top of a plane inclined at an angle 45° with the horizontal. How far from the point of projection with the ball strike the plane?


- (a) $\frac{v^2}{g}$ (b) $\sqrt{2} \frac{v^2}{g}$ (c) $\frac{2v^2}{g}$ (d) $\sqrt{2} \left[\frac{2v^2}{g} \right]$

Q 11. Two inclined planes OA and OB having inclination with horizontal 30° and 60° respectively, intersect each other at O as shown in figure. A particle is projected from point P with velocity $u = 10\sqrt{3}$ m/s Along a direction perpendicular to plane OA. If the particle strikes plane OB perpendicularly at Q, calculate Velocity with which particle strikes the plane OB? ($g = 10$ m/s²)



- (a) 10 m/s (b) $10\sqrt{3}$ m/s
 (c) $\sqrt{3}$ m/s (d) $5\sqrt{3}$ m/s


Q 12. A particle P is projected from a point on the surface of smooth inclined plane (see figure). Simultaneously another particle Q is released on the smooth inclined plane from the same position. P and Q collide after $t = 4$ second. The speed of projection of P is :-





- (a) 5 m/s (b) 10 m/s
(c) 15 m/s (d) 20 m/s

Q 13. A ball is projected on smooth inclined plane in direction perpendicular to line of greatest slope with velocity of 8m/s. Find it's speed after 1 sec.



- (a) 10 m/s (b) 12 m/s
(c) 15 m/s (d) 20 m/s

Q 14. A particle is projected from a point P(2,0,0)m with a velocity 10m/s making an angle 45° with the horizontal. The plane of projectile motion passes through a horizontal line PQ which makes an angle of 37° with positive x-axis, xy plane is horizontal. The coordinates of the point where the particle will strike the line PQ is : -(take $g = 10 \text{ m/s}^2$)

- (a) (10,6,0)m (b) (8,6,0)m (c) (10,8,0)m (d) (6,10,0)m

Answer Key

Q.1 b	Q.2 d	Q.3 a	Q.4 a	Q.5 a
Q.6 c	Q.7 d	Q.8 a	Q.9 b	Q.10 d
Q.11 a	Q.12 b	Q.13 a	Q.14 a	


PLUS **ICONIC****

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months	₹2,333/mo	>
No cost EMI	₹56,000	
18 months	₹2,625/mo	>
No cost EMI	₹47,250	
12 months	₹3,208/mo	>
No cost EMI	₹38,500	
6 months	₹4,667/mo	>
No cost EMI	₹28,000	

To be paid as a one-time payment

[View all plans](#)

 Add a referral code APPLY

PHYSICSLIVE


PLUS **ICONIC****

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months	₹2,100/mo	>
No cost EMI	+10% OFF ₹50,400	
18 months	₹2,363/mo	>
No cost EMI	+10% OFF ₹42,525	
12 months	₹2,888/mo	>
No cost EMI	+10% OFF ₹34,650	
6 months	₹4,200/mo	>
No cost EMI	+10% OFF ₹25,200	

To be paid as a one-time payment

[View all plans](#)

 Awesome! **PHYSICSLIVE** code applied ✗

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS.

Written Solution

DPP-6 Projectile Motion on inclined plane

By Physicsaholics Team

Q) On an inclined plane of inclination 30° , a ball is thrown at an angle of 60° with the horizontal from the foot of the incline with a velocity of $10\sqrt{3}$ m/s. If $g = 10$ m/s², then find the time in which ball will hit the inclined plane?

(a) 1 s

(b) 2 s

(c) 3 s


(d) 4 s

PRATEEK JAIN
PHYSICSAHOLICS



Ans. b

Solution:



$$T = \frac{2u_y}{g}$$
$$= \frac{2 \times \frac{10\sqrt{3}}{2}}{\frac{10\sqrt{3}}{2}}$$
$$= 2 \text{ Sec}$$

Q) Two bodies are projected from the same point with equal speeds in such a directions that they strike on the same point on a plane whose inclination is β . If α the angle of projection of the first, ratio of there times of flight is

(a) $\frac{\sin \alpha}{\cos \beta}$

(b) $\frac{\sin \alpha}{\sin \beta}$

(c) $\frac{\sin(\alpha - \beta)}{\cos \beta}$



(d) $\frac{\sin(\alpha - \beta)}{\cos \alpha}$

PRATEEK JAIN
PHYSICSAHOLICS



Ans. d

Solution:



at angle α

$$T_1 = \frac{2u \sin(\alpha - \beta)}{g \cos \beta}$$

at angle α'

$$T_2 = \frac{2u \sin(\alpha' - \beta)}{g \cos \beta}$$


$$\frac{T_1}{T_2} = \frac{\sin(\alpha - \beta)}{\sin(90 - \alpha)} = \frac{\sin(\alpha - \beta)}{\cos \alpha}$$

by symmetry
 $\alpha - \beta = 90 - \alpha'$

$$\alpha' = 90 - (\alpha - \beta)$$


Q) A baseball is projected with a velocity v making an angle θ with the incline of inclination α as shown in fig. Find the condition that the ball hits the incline at right angle.

- (a) $\cot \theta = 2 \tan \alpha$ (b) $\sin \theta = \cos \alpha$
(c) $\tan \theta = \sin \alpha$ (d) $\cot \theta = \cos \alpha$



Ans. a

Solution:



time of flight $T = \frac{2u_y}{g_y}$

$$T = \frac{2v \sin \theta}{g \cos \alpha}$$

at $t = T$, $v_x = 0$

but $v_x = u_x + a_x t$

$$\Rightarrow 0 = v \cos \theta - g \sin \alpha \times \frac{2v \sin \theta}{g \cos \alpha}$$

$$\Rightarrow \cos \theta = \frac{2 \sin \alpha \sin \theta}{\cos \alpha}$$

$$\Rightarrow \cot \theta = 2 \tan \alpha$$

Q) A projectile is required to hit a target whose coordinates relative to horizontal and vertical axes through the point of projection are (α, β) . If the gun velocity is $\sqrt{2g\alpha}$, it is impossible to hit the target if

(a) $\beta > 3\alpha/4$

(b) $\beta \geq 1\alpha/4$

(c) $\beta \leq 3\alpha/4$

(d) $\beta \geq 3\alpha/4$

PRATHEEK JAIN
PHYSICSAHOLICS



Ans.


Solution:

Maximum Range on
Inclined plane

$$R_{\max} = \frac{u^2}{g(1 + \sin \theta)}$$

$$= \frac{2gd}{g\left(1 + \frac{\beta}{\sqrt{d^2 + \beta^2}}\right)}$$

If it is not possible to hit target
If $R_{\max} < \sqrt{d^2 + \beta^2}$



$$\frac{2d}{1 + \frac{\beta}{\sqrt{d^2 + \beta^2}}} < \sqrt{d^2 + \beta^2}$$

$$\Rightarrow 2d < \sqrt{d^2 + \beta^2} + \beta$$

$$\Rightarrow 2d - \beta < \sqrt{d^2 + \beta^2}$$

$$\Rightarrow 4d^2 + \beta^2 - 4d\beta < d^2 + \beta^2$$

$$\Rightarrow 3d^2 < 4d\beta \Rightarrow \beta > \frac{3d}{4}$$

Q) A marble rolls down from top of a staircase with constant horizontal velocity 10 m/s. If each step is $y = 1$ meter high and $x = 1$ meter wide. To which step the marble will strike directly? ($g = 9.8 \text{ m/s}^2$)

(a) 21st

(b) 8th


(c) 10th

(d) 18th



Ans. a

Solution:



Range of ball on inclined plane
 $R = \frac{2u^2 \cos(45^\circ - 45^\circ) \sin 45^\circ}{g \cos^2 45^\circ}$

$$R = \frac{2 \times 100 \times 1 \times \frac{1}{\sqrt{2}}}{g \times \frac{1}{2}}$$

$$= \frac{200 \sqrt{2}}{g} = (20 \dots) \sqrt{2}$$

Since $20\sqrt{2} < R < 21\sqrt{2}$

\Rightarrow ball will hit 21st step

\downarrow
Imaginary inclined plane

If $R < \sqrt{2}$
 \Rightarrow ball will hit first step.
If $\sqrt{2} < R < 2\sqrt{2}$
 \Rightarrow ball will hit 2nd step

Q) A particle is projected from origin of coordinate system. A target is fixed at point (40m, 30m). Find the minimum velocity of projectile to hit the target? ($g = 10 \text{ m/s}^2$)

(a) 10 m/s

(b) 17 m/s

(c) $20\sqrt{2} \text{ m/s}$

(d) $10\sqrt{5} \text{ m/s}$

PRATEEK JAIN
PHYSICSAHOLICS



Ans. c

Solution:


$$R_{\max} = \frac{u^2}{g(1 + \sin \alpha)}$$

$$\Rightarrow 50 = \frac{u^2}{10(1 + \frac{30}{50})}$$

$$\Rightarrow u^2 = 800 \Rightarrow u = 20\sqrt{2} \text{ m/sec.}$$

★ at $u = 20\sqrt{2}$ max Range is 50m. means it is just possible to hit target.

If $u < 20\sqrt{2}$ max Range will be less than 50m & particle will not be able to hit target.




Q) A particle is projected at an angle θ with an inclined plane making an angle β with the horizontal as shown in figure, speed of the particle is u , after time t find y component of velocity when particle is at maximum distance from the incline plane ?

(a) $\frac{u}{\sqrt{2}}$

(b) $\frac{2u}{\sqrt{3}}$


(c) $\frac{\sqrt{2}u}{3}$

(d) zero



Ans. d

Solution:



Projection of motion on y axis \rightarrow

$$v_y = 0$$

$$u_y$$

at maximum distance from

Incline $v_y = 0$


Q) A ball is projected from point A with a velocity 10 m/s perpendicular to the inclined plane as shown in figure. Range of the ball on the inclined plane is :

(a) $\frac{40}{3}$ m

(c) $\frac{13}{20}$ m


(b) $\frac{20}{13}$ m

(d) $\frac{13}{40}$ m



Ans. a

Solution:



$$T = \frac{2u_y}{g_y} = \frac{2 \times 10}{g\sqrt{3}/2} = \frac{4}{\sqrt{3}}$$

$$R = u_x T + \frac{1}{2} a_x T^2$$
$$= 0 + \frac{1}{2} \times \frac{g}{2} \times \frac{16}{3}$$
$$= \frac{40}{3} \text{ m}$$

Q) A plane surface is inclined making an angle θ with the horizontal. From the bottom of this inclined plane, a bullet is fired with velocity v . The maximum possible range of the bullet on the inclined plane is

(a) $\frac{v^2}{g}$

(b) $\frac{v^2}{g(1+\sin \theta)}$


(c) $\frac{v^2}{g(1-\sin \theta)}$

(d) $\frac{v^2}{g(1+\cos \theta)}$



Ans. b

Solution:



$$R = \frac{2u^2 \cos(\theta + \alpha) \sin \alpha}{g \cos^2 \theta}$$

$$R = \frac{u^2 [\sin(2\theta + \alpha) - \sin \alpha]}{g(1 - \sin^2 \theta)}$$

$$R_{\max} = \frac{u^2 (1 - \sin \alpha)}{g(1 - \sin \alpha)(1 + \sin \alpha)}$$
$$= \frac{u^2}{g(1 + \sin \alpha)}$$

Q) A ball is projected horizontal with a speed v from the top of a plane inclined at an angle 45° with the horizontal. How far from the point of projection with the ball strike the plane?

(a) $\frac{v^2}{g}$

(b) $\sqrt{2} \frac{v^2}{g}$

(c) $\frac{2v^2}{g}$


(d) $\sqrt{2} \left[\frac{2v^2}{g} \right]$

PRATEEK JAIN
PHYSICSAHOLICS



Ans. d

Solution:





$$T = \frac{2U_y}{g} = \frac{2v/\sqrt{2}}{g/\sqrt{2}} = \frac{2v}{g}$$

$$R = U_x T + \frac{1}{2} a_x T^2$$
$$= \frac{v}{\sqrt{2}} \times \frac{2v}{g} + \frac{1}{2} \times \frac{g}{\sqrt{2}} \frac{4v^2}{g^2}$$
$$= \frac{4v^2}{g\sqrt{2}} = \frac{2\sqrt{2}v^2}{g}$$

Q) Two inclined planes OA and OB having inclination with horizontal) 30° and 60° respectively, intersect each other at O as shown in figure. A particle is projected from point P with velocity $u = 10\sqrt{3}$ m/s Along a direction perpendicular to plane OA. If the particle strikes plane OB perpendicularly at Q, calculate Velocity with which particle strikes the plane OB? ($g = 10$ m/s²)


- (a) 10 m/s
(c) $\sqrt{3}$ m/s

- (b) $10\sqrt{3}$ m/s
(d) $5\sqrt{3}$ m/s



Ans. a

Solution:



for motion along x axis

$$V_x = u_x + a_x t$$

$$0 = 10\sqrt{3} - \frac{g\sqrt{3}}{2} t$$

$$t = 2 \text{ Sec}$$

for motion along y axis

$$V_y = u_y + a_y t = 0 + \frac{g}{2} \times 2$$

$$= 10 \text{ m/sec} \quad \text{Ans} = 10 \text{ m/sec}$$


Q) A particle P is projected from a point on the surface of smooth inclined plane (see figure). Simultaneously another particle Q is released on the smooth inclined plane from the same position. P and Q collide after $t = 4$ second. The speed of projection of P is :-

(a) 5 m/s

(b) 10 m/s


(c) 15 m/s

(d) 20 m/s



Ans. b

Solution:



for motion along x axis
P & Q both have same
displacement, same time
of motion & same
acceleration ($g \sin 60^\circ$)

Therefore their u_x must be same $\Rightarrow u_x = 0$
for P, $T = t = \frac{2u_y}{g} \Rightarrow \frac{2u_y}{g \cos 60^\circ} = t$

$u_y = 10 \text{ m/sec} \Rightarrow$ initial velocity of P = 10 m/sec


Q) A ball is projected on smooth inclined plane in direction perpendicular to line of greatest slope with velocity of 8m/s . Find it's speed after 1 sec.

(a) 10 m/s

(b) 12 m/s


(c) 15 m/s

(d) 20 m/s



Ans. a

Solution:



motion along x axis is uniform

$$\Rightarrow V_x = 8 \text{ m/sec}$$

motion along y axis has acceleration 6 m/sec^2 .

$$V_y = U_y + a_y t = 0 + 6 \times 1 = 6 \text{ m/sec}^2$$

$$g \sin 37^\circ = 10 \times \frac{3}{5} = 6 \text{ m/sec}^2$$

$$\underline{V} = 8\hat{i} + 6\hat{j}$$
$$V = 10 \text{ m/sec}$$

Q) A particle is projected from a point P(2,0,0)m with a velocity 10m/s making an angle 45° with the horizontal. The plane of projectile motion passes through a horizontal line PQ which makes an angle of 37° with positive x-axis, xy plane is horizontal. The coordinates of the point where the particle will strike the line PQ is : -(take $g = 10 \text{ m/s}^2$)

(a) (10,6,0)m

(b) (8,6,0)m


(c) (10,8,0)m

(d) (6,10,0)m



Ans. a

Solution:



$$\text{Range} = \frac{u^2 \sin 2\theta}{g}$$
$$= \frac{10 \times 10}{10} = 10\text{m}$$

Co-ordinates of point of fall
 $= (10, 6, 0)$

For Video Solution of this DPP, Click on below link

Video Solution
on Website:-

<https://physicsaholics.com/home/courseDetails/52>

Video Solution
on YouTube:-

https://youtu.be/F0BVSJhp_k

Written Solution
on Website:-

<https://physicsaholics.com/note/notesDetails/74>

 **SUBSCRIBE**



[@Physicsaholics](#)

[@Physicsaholics_prateek](#)

[@NEET_Physics](#)
[@IITJEE_Physics](#)

[physicsaholics.com](#)

[Unacademy](#)



CLICK

Chalo Niklo