

Video Solution on Website:-

<https://physicsaholics.com/home/courseDetails/53>


Video Solution on YouTube:-

<https://youtu.be/68f925ejomw>


Written Solution on Website:-

<https://physicsaholics.com/note/notesDetails/75>


- Q 1. A frictionless cart of mass  $M$  carries two other frictionless carts having masses  $m_1$  and  $m_2$  connected by a string passing over a pulley as shown in figure. The horizontal force that must be applied on  $M$  so that  $m_1$  and  $m_2$  do not move relative to it will be -



- (a)  $(M + m_1 + m_2) (m_2 / m_1) g$   
 (b)  $(M + m_1 + m_2) (m_1 / m_2) g$   
 (c)  $(M + m_1) [(m_1 + m_2) / m_2] g$   
 (d)  $(M + m_2) [m_2 / (m_1 + m_2)] g$
- Q 2. Two cubes of masses  $m_1$  and  $m_2$  lie on frictionless slopes of a block A which rests on a horizontal table. The cubes are connected by a string which passes over a pulley as shown in figure. If  $a_0$  be the horizontal acceleration to which the whole system (block + masses) is subjected so that  $m_1$  and  $m_2$  do not move and  $T$  be the tension in the string in that situation then-




- (a)  $a_0 = \left( \frac{m_1 \sin \alpha + m_2 \sin \beta}{m_1 + m_2} \right) g$   
 (b)  $a_0 = \left( \frac{m_1 \sin \alpha + m_2 \sin \beta}{m_1 \cos \alpha + m_2 \cos \beta} \right) g$   
 (c)  $T = \frac{m_1 m_2}{m_1 + m_2} g \sin(\alpha + \beta)$   
 (d)  $T = \left( \frac{m_1 m_2}{m_1 \cos \alpha + m_2 \cos \beta} \right) g \sin(\alpha - \beta)$
- Q 3. In the arrangement shown in figure wedge of mass  $M$  moves towards left with an acceleration  $a$ . All surfaces are smooth. The acceleration of block in relative to wedge is:




- (a)  $a/2$                       (b)  $\frac{2Ma}{m}$   
 (c)  $\frac{a}{2} + \frac{g\sqrt{3}}{2}$                 (d)  $\frac{(M+m)a}{m}$

Q 4. A block is placed in a smooth cylinder which is moving horizontally with constant acceleration  $a = 3g/4$ . Find height of block from bottommost point of cylinder if block is stationary with respect to cylinder ?




- (a)  $R/5$                       (b)  $R/3$   
 (c)  $R/4$                       (d)  $R/2$

Q 5. x-y plane is a vertical plane in which a parabolic wire of shape  $y = x^2$  is moving with constant acceleration  $a$  in negative x direction. At position shown in figure a bead is stationary with respect to wire. Find height of bead ?




- (a)  $a/g$                       (b)  $a/2g$   
 (c)  $\frac{a^2}{4g^2}$                       (d) none of these

Q 6. In given figure all surfaces are smooth and string is massless. System is released from given position. Find initial acceleration of cart ?




- (a)  $g/7$                       (b)  $2g/7$   
 (c)  $\frac{g\sqrt{3}}{7}$                       (d)  $\frac{g}{7\sqrt{3}}$

Q 7. After releasing triangular wedge of mass  $m$  moves left by acceleration  $g/10$ . find mass of block if all surfaces are smooth ?



- (a)  $m$                                       (b)  $m/37$   
 (c)  $5m/52$                               (d)  $25m/104$

Q 8. A pulley fixed to the ceiling of an elevator car carries a thread whose ends are attached to the masses  $m_1 = 3 \text{ kg}$  and  $m_2 = 6 \text{ kg}$ . The car starts going up with an acceleration  $a_0 = 2 \text{ m/sec}^2$ . Assuming the masses of the pulley and the thread as well as the friction to be negligible, find acceleration of  $m_1$  with respect to ground ?



- (a)  $4 \text{ m/Sec}^2$                       (b)  $2 \text{ m/Sec}^2$   
 (c)  $6 \text{ m/Sec}^2$                       (d)  $8 \text{ m/Sec}^2$

## Answer Key

<b>Q.1 a</b>	<b>Q.2 b,d</b>	<b>Q.3 c</b>	<b>Q.4 a</b>	<b>Q.5 c</b>
<b>Q.6 c</b>	<b>Q.7 d</b>	<b>Q.8 c</b>		



## NEET UG subscription

PLUS

ICONIC\*\*

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months ₹2,100/mo >  
No cost EMI +10% OFF ₹50,400

18 months ₹2,363/mo >  
No cost EMI +10% OFF ₹42,525

12 months ₹2,888/mo >  
No cost EMI +10% OFF ₹34,650

6 months ₹4,200/mo >  
No cost EMI +10% OFF ₹25,200

To be paid as a one-time payment

[View all plans](#)



Awesome! **PHYSICSLIVE** code applied



# PHYSICSLIVE

Use code **PHYSICSLIVE** to get 10% OFF on Unacademy PLUS and learn from India's Top Faculties.



## NEET UG subscription

PLUS

ICONIC\*\*

- ✓ India's Best Educators
- ✓ Interactive Live Classes
- ✓ Structured Courses & PDFs
- ✓ Live Tests & Quizzes
- ✗ Personal Coach
- ✗ Study Planner

24 months ₹2,100/mo >  
No cost EMI +10% OFF ₹50,400

18 months ₹2,363/mo >  
No cost EMI +10% OFF ₹42,525

12 months ₹2,888/mo >  
No cost EMI +10% OFF ₹34,650

6 months ₹4,200/mo >  
No cost EMI +10% OFF ₹25,200

To be paid as a one-time payment

[View all plans](#)



Awesome! **PHYSICSLIVE** code applied



# Written Solution

**DPP-6 NLM: Pseudo Force**


**By Physicsaholics Team**



## Solution:1

Let acceleration  
of  $M$  is  $a$ .

$$F = (M + m_1 + m_2)a \quad \text{--- (i)}$$



Since  $m_1$  &  $m_2$  are not sliding on  $M$ .

Supporting force = opposing force

$$\Rightarrow m_1 a = m_2 g \Rightarrow a = \frac{m_2 g}{m_1}$$

$$\Rightarrow F = (m_1 + m_2 + M) \frac{m_2 g}{m_1}$$

Ans. (A)

## Solution:2


Since blocks are not sliding.

Supporting forces  
= opposing forces

$$\Rightarrow m_1 g \sin \alpha + m_2 g \sin \beta = m_1 a_0 \cos \alpha + m_2 a_0 \cos \beta$$

$$\Rightarrow a_0 [m_1 \cos \alpha + m_2 \cos \beta] = (m_1 \sin \alpha + m_2 \sin \beta) g$$

$$\Rightarrow a_0 = \frac{m_1 \sin \alpha + m_2 \sin \beta}{m_1 \cos \alpha + m_2 \cos \beta} g$$



Ans.b,d

Solution.3

from F.B.D of  $m_1$

$$T + m_1 a_0 \cos \alpha = m_1 g \sin \alpha$$

on putting value of  $a_0$

$$T = \frac{m_1 m_2 g \sin(\alpha - \beta)}{m_1 \cos \alpha + m_2 \cos \beta}$$


Ans. (B, D)



Solution: 4

$$\mu a \cos 60^\circ + mg \sin 60^\circ = \mu a_0$$

$$a_0 = \frac{a}{2} + \frac{g\sqrt{3}}{2}$$



Ans. (c)

Solution:5


Resultant of  $ma$   
&  $mg$  must be  
just opposite to  
 $N$ .

$$\tan \theta = \frac{ma}{mg}$$

$$\Rightarrow \tan \theta = \frac{3}{4}$$

$$\Rightarrow \cos \theta = \frac{4}{5} = \frac{R-h}{R}$$

$$\Rightarrow h = R/5$$



Ans. (A)


## Solution:6

Resultant of  $ma$   
&  $mg$  must be  
perpendicular to  $N$ .

$$\Rightarrow \tan \theta = \frac{ma}{mg}$$

$$\Rightarrow \frac{dy}{dx} = \frac{a}{g} \Rightarrow 2x = \frac{a}{g}$$

$$\Rightarrow y = x^2 = \frac{a^2}{4g^2}$$



Ans.c



Solution:7

for Cart

$$T \sin 60^\circ = ma$$


$$\Rightarrow T = \frac{2ma}{\sqrt{3}}$$

forces along string on bob must be balanced.

$$T + ma \cos 30^\circ = mg \cos 60^\circ$$


$$\frac{2ma}{\sqrt{3}} + \frac{ma\sqrt{3}}{2} = \frac{mg}{2}$$

$$\Rightarrow a = \frac{g\sqrt{3}}{7}$$



Ans.c

## Solution:8



Ans.d



Solution.9


forces along normal  
must be balanced.

$$\Rightarrow N + \frac{m_1 g}{10} \times \frac{4}{5}$$

$$= m_1 g \times \frac{3}{5}$$

$$\Rightarrow \frac{m_1 g}{8} = m_1 g \left( \frac{3}{5} - \frac{2}{25} \right)$$

$$\Rightarrow m_1 = \frac{25g}{104}$$



Ans. (d)


Solution:10

$$g_{\text{eff}} = g + a_0 = 10 + 2 \\ = 12 \text{ m/Sec}^2$$

$$a = \frac{m_1 - m_2}{m_1 + m_2} g_{\text{eff}} \\ = \frac{6 - 3}{6 + 3} \times 12 = 4 \text{ m/Sec}^2$$

Here  $a$  is acceleration of blocks w.r.t. lift

$$a_{m,g} = a_{m,L} + a_{L,g} = 4 \uparrow + 2 \uparrow \\ = 6 \text{ m/Sec}^2$$



Ans. (C)

For Video Solution of this DPP, Click on below link

Video Solution  
on Website:-

<https://physicsaholics.com/home/courseDetails/53>

Video Solution  
on YouTube:-

<https://youtu.be/68f925ejomw>

Written Solution  
on Website:-

<https://physicsaholics.com/note/notesDetails/75>

 **SUBSCRIBE**



[@Physicsaholics](#)

[@Physicsaholics\\_prateek](#)

[@NEET\\_Physics](#)

[@IITJEE-Physics](#)

[physicsaholics.com](#)

[Unacademy](#)



**CLICK**

Chalo Niklo